arXiv:1212.0051v1 [math.GT] 1 Dec 2012

Prescribing the behavior of Weil-Petersson geodesics in the moduli space of Riemann surfaces

Babak Modami

Abstract

In this paper we study the Weil-Petersson (WP) geodesics with *narrow ending data* and develop techniques to control length-functions and twist parameters along them and prescribe their itinerary in the moduli space. This class of geodesics is rich enough to provide for examples of closed WP geodesics in the thin part of moduli space, as well as diverging WP geodesic rays with minimal filling ending lamination.

We show that the strong asymptotic class of rays with narrow ending data and bounded annular coefficients is determined by the forward ending data, generalizing the *Recurrent ending lamination theorem* of Brock, Masur and Minsky for WP geodesic rays recurrent to the thick part of moduli space.

As an intermediate step we characterize all the stable hierarchy resolution paths in the pants graph of the surface.

Contents

1 Introduction **2**

2 Curve complexes and hierarchy machine **7**
2.1 The pants and marking graph 10
2.2 Hierarchies and their resolutions in the pants and marking graph 12
2.3 $\Sigma-$hulls and their projections 14

3 The Weil-Petersson metric and its synthetic properties **15**
3.1 Ending data 19

4 Compactness arguments in the Weil-Petersson completion of Teichmüller space **20**
4.1 Limits of sequences of uniformly bounded length WP geodesic segments 21
4.2 Length-function versus twist parameter control 25

5 Stable hierarchy paths **30**
5.1 Narrow ending data 31
5.2 Stability in the pants graph 36

5.3 Fellow-traveling 45

6 Itinerary of a Weil-Petersson geodesic segment 46
6.1 Isolated annular subsurfaces 47
6.2 Length-function control 50

7 Weil-Petersson Geodesics 59
7.1 Weil-Petersson geodesic rays with prescribed itinerary 59
7.2 Diverging geodesic rays 62
7.3 Closed geodesics in the thin part 63
7.4 A recurrence criterion 64

8 Asymptotic class of rays with narrow ending data 64
8.1 Bounding annular coefficients 65
8.2 Strongly asymptotic rays 67

9 Appendix : Ending data with prescribed subsurface coefficients 75

1 Introduction

The Weil-Petersson (WP) metric on the moduli space of Riemann surfaces is an incomplete Riemannian metric with sectional curvatures asymptotic to both 0 and $-\infty$ in the completion. It is a metric of higher rank in the sense of Gromov, see [BF06]. These features prevent applying most of the standard techniques available in the study of the global geometry and dynamics of complete metrics with negative curvature bounded away from zero and rank 1 metrics (for example see [PP10], [Ebe72]) to the WP metric. The main theme of this paper and the previous work of Brock, Masur and Minsky (see [BM08][BMM10][BMM11]) is to apply combinatorial techniques from low dimensional topology to study the global behavior of WP geodesics.

In [BMM10] the authors introduce a notion of ending lamination for WP geodesic rays. They show that the ending lamination determines the strong asymptotic class of a WP geodesic ray recurrent to a compact subset of moduli space. They show that there are WP geodesics dense in the moduli space. In [BMM11], a more explicit connection between the combinatorics of the ending laminations of a WP geodesic and its behavior was established, using which they get estimates on the number of closed geodesics of a bounded length in compact subsets of moduli space. In this paper we prove the following two results about the behavior of WP geodesics:

Theorem 1.1. *Given any compact subset of moduli space $\mathcal{K}$ there are closed Weil-Petersson geodesics not intersecting $\mathcal{K}$.*

Theorem 1.2. *Starting from any point in the moduli space there are uncountably many diverging WP geodesic rays with minimal filling ending lamination.*

By the Poincaré recurrence theorem almost every WP geodesic ray is recurrent to a compact subset of moduli space, however the above theorem exhibits the abundance of WP geodesic rays diverging in the moduli space in an interesting way, more precisely these geodesics diverge in the moduli space by getting closer and closer to a chain of completion strata.

Given a WP geodesic $g : (a, b) \to \mathrm{Teich}(S)$ denote the ending data associated to its forward trajectory $g|_{[0,b)}$ by $\nu^+ = \nu^+(g)$ and the one associated to its backward trajectory $g|_{[0,a)}$ by $\nu^- = \nu^-(g)$, each of $\nu^\pm$ is either a lamination or a partial marking.

To each subsurface $Y \subseteq S$ that is not a three holed sphere, there is an associated *subsurface coefficient* denoted by

$$d_Y(\nu^-, \nu^+)$$

which is the distance in the curve complex of Y between the projections of ν^- and ν^+, for more details see §2.

An $A-$*narrow* condition on the ending data is a constraint on the set of subsurfaces with large subsurface coefficient, more precisely a pair (ν^-, ν^+) is $A-$narrow if every non-annular $Y \subseteq S$ with

$$d_Y(\nu^-, \nu^+) > A$$

is *non-separated* i.e. is a connected subsurface with complement consisting of only annuli and three holed spheres.

We prove the following generalization of the *Ending lamination theorem* of recurrent WP geodesic rays in [BMM10]:

Theorem 1.3. *(Narrow ending lamination theorem) The strong asymptotic class of a WP geodesic ray with narrow ending data and bounded annular coefficients is determined by the forward ending lamination* ν^+*.*

The class of geodesic rays with narrow ending data and bounded annular coefficients contain rays which are not recurrent to any compact subset of moduli space (diverging rays) as well. Heuristically these geodesic rays avoid quasi-flats in the WP metric and exhibit features of geodesic in spaces of negative curvature.

Subsurface coefficients are an analogue of continued fraction expansions which provide for a coding of geodesics on the modular surface, see for example [Ser85]. Conjecturally these coefficients give extensive information about the behavior of Weil-Petersson geodesics in the moduli space. The following conjecture was proposed in [BMM10]:

Conjecture 1.4. (Short curves) Let g be a Weil-Petersson geodesic with ending data (ν^-, ν^+). Then

1. For every $\epsilon > 0$ there is $A > 0$, such that if $d_Y(\nu^-, \nu^+) > A$ then $\inf_t \ell_\alpha(g(t)) \leq \epsilon$ for every $\alpha \in \partial Y$.

2. For every $A > 0$ there is $\epsilon > 0$, such that if $\inf_t \ell_\alpha(g(t)) \leq \epsilon$ then there is $Y \subsetneq S$ such that $\alpha \in \partial Y$ and $d_Y(\nu^-, \nu^+) > A$.

Also one would hope for estimates on the amount of time that a curve is short (has length less than a given ϵ) along a WP geodesic.

The WP metric exhibits different features in the thick and thin parts of Teichmüller space, for instance in any thick part the sectional curvatures are all bounded away from both 0 and $-\infty$, while in the thin parts the WP metric is almost a product metric. So many questions in the study of the dynamics of the WP metric reduce to determining the itinerary of geodesics, that is given a WP geodesic g one would hope to predict the thick and thin parts of Teichmüller space where g visits, the time that g spends in each, and the order in which g visits them.

After the work of Masur-Minsky and Rafi (see [MM99] [Raf05][Raf07] and [Mas09] for a survey) the above conjecture holds for Teichmüller geodesics and provides for a complete picture of the itinerary of Teichmüller geodesics.

The underlying machinery of these results is the Masur-Minsky machinery of hierarchies and their resolutions in pants and marking graphs introduced in [MM00]. Given a pair of partial markings or laminations (ending data) a *hierarchy (resolution) paths* $\rho : [m,n] \to P(S)$ ($[m,n] \subseteq \mathbb{Z}$) between them is a quasi-geodesic in the pants graph with certain properties encoded in the ending data and their subsurface coefficients. For example corresponding to any subsurface Z with large enough coefficient $d_Z(\nu^-, \nu^+)$ there are subintervals of $[m,n]$ denoted by J_Z, see Theorem 2.11 for the list of properties.

Brock's quasi-isometry, Theorem 3.3, asserts that the Bers pants decompositions along a WP geodesic trace a quasi-geodesic in the pants graph of the surface. When such a quasi-geodesic fellow-travels (see Definition 5.14) a hierarchy path ρ we have a correspondence between the parameters along the hierarchy path and that of the WP geodesic, which is roughly the nearest point correspondence of fellow-traveling paths. In this situation one would hope that the hierarchy machinery helps to determine the itinerary of WP geodesics. An example would be visiting the thin part of a multi-curve σ over the time interval corresponding to J_σ.

The $A-$narrow condition on ending data turns out to imply a uniform fellow-traveling of a WP geodesics segment and a hierarchy paths with the same ending data. In this paper we develop a control for length-functions and twist parameters along these geodesics and show that their itinerary mimic combinatorial properties of hierarchy paths. In order to prove our main technical results we introduce the following notion:

Let $Z \subseteq S$ be a subsurface with large enough subsurface coefficient then we say that Z has $(R, R')-$*bounded combinatorics* over a subinterval $[i_1, i_2] \subset J_Z \subset [m,n]$ if for every non-annular subsurface $Y \subsetneq Z$

$$d_Y(\rho(i_1), \rho(i_2)) \leq R$$

and for every annular subsurface $A(\gamma)$ with core curve γ inside Z

$$d_\gamma(\rho(i_1), \rho(i_2)) \leq R'$$

This condition is a local version of the bounded combinatorics of the ending data in [BMM11] which implies that the geodesic stays in the thick part of moduli space.

We say that a subsurface Z has $(R,R')-$bounded combinatorics over a subinterval along a WP geodesic if the geodesic fellow-travels a subinterval of a hierarchy path where Z has $(R,R')-$bounded combinatorics. In the direction of conjecture 1.4 we prove:

Theorem 1.5. *Given positive constants A, R and R' and $\epsilon > 0$ sufficiently small, there are constants $\bar{s}(A,R,R',\epsilon)$ and $\bar{\epsilon}(A,R)$ with the following property:*

Let $g : [a,b] \to \mathrm{Teich}(S)$ be a Weil-Petrsson geodesic segment parametrized by arc-length with $A-$narrow ending data. Let a non-separated subsurface $Z \subseteq S$ have $(R,R')-$bounded combinatorics over $[t_1,t_2] \subset [a,b]$.

If $t_2 - t_1 > 2\bar{s}$ then

1. *$\ell_\alpha(g(t)) \leq \epsilon$ for any $\alpha \in \partial Z$*
2. *$\ell_\gamma(g(t)) \geq \bar{\epsilon}$ for every $\gamma \notin \partial Z$*

for every $t \in [t_1 + \bar{s}, t_2 - \bar{s}]$.

In §6 we use bounded combinatorics interval to isolate annular subsurface along a hierarchy path. The twist parameter developed about an *isolated annular subsurface* along a WP geodesic is comparable to the one along a fellow-traveling hierarchy path. This together with the length-function versus twist parameter controls over uniformly bounded length WP geodesics segments we develop in §4 are main technical tools in §6 where we prove the above theorem.

Using the control we develop on length-functions along WP geodesic segments and by extracting limits of WP geodesic segments with narrow ending data in Theorem 7.2 we provide WP geodesics with prescribed itinerary in the moduli space. Itineraries of these rays mimic the combinatorial properties of hierarchy paths encoded in the ending data. Theorems 1.1 and 1.2 will follow from our constructions in section 7. For similar examples in $\mathcal{M}(S_{1,1})$ see [PWW10]. Also see the work of Jeff Brock in [Bro].

In our setting fellow-traveling is part of the combinatorial frame work to control length-functions. In §5 we give the following characterization of stable hierarchy paths in the pants graph of a surface:

Theorem 5.8. *(Stable hierarchy path) Given $A > 0$ there is a function D such that any $A-$narrow hierarchy path is $D-$stable in the pants graph. Moreover, given D there is $A' > 0$ such that if a hierarchy path is $D-$stable then it is A'-narrow.*

In this theorem $D : \mathbb{R}^{\geq 1} \times \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ is the quantifier of the stability, see Definition 5.1.

This theorem implies that the Bers pants along a WP geodesic and a hierarchy path with the same narrow ending data uniformly fellow-travel each other in the pants graph.

Theorem 1.3 follows from our control on length-functions along WP geodesics together with some ruled surface arguments as in [BMM10]. The main new ingredient in our proof is a kind of *geometric limit* argument which guarantees the existence of an infinite sequence of regions with a definite negative total curvature on ruled surfaces with one side on a ray with prescribed itinerary. This result helps us to overcome some of the difficulties due to the fact that the curvatures of the WP metric are not bounded away from 0 and prove visibility of the class of geodesic rays with narrow ending data and bounded annular coefficients.

Outline of the paper:

Section 2 is devoted to background about curve complexes and some important notions and results in the setting of pants and marking graphs. We also recall hierarchical structures on pants and marking graphs and their resolutions introduced by Masur and Minsky in [MM00]. In Theorem 2.11 we list the properties of hierarchy resolution paths. In this section we also recall the $\Sigma-$hulls and their projections from [BKMM12].

In section 3 we give some background about the WP metric and synthetic properties of WP geodesics.

In section 4 we provide the proofs of refined version of some of the key results in [BMM11] and [Wol03] which we need in this work. Some of them were suggested to us by Jeffery Brock. These are mainly based on compactness arguments in the setting of the WP metric and are consequences of Wolpert's geodesic limit theorem. These results give us a kind of control on development of Dehn twists versus length-functions along uniformly bounded length WP geodesic segments.

In section 5 we give a characterization of stable hierarchy paths. These are hierarchy paths with narrow ending data. The proof will be through the $\Sigma-$hulls and their projections.

In section 6, which is the heart of our paper, we develop some new techniques to control the length-functions and twist parameters along WP geodesics fellow-traveling a hierarchy paths, see Lemma 6.8 and Theorem 6.1. The control is over a subinterval over which a WP geodesic fellow-travels a subinterval of a hierarchy path over which all annular subsurfaces except possibly a few of them which consist a multi-curve are isolated.

In section 7 we provide examples of WP geodesics with prescribed behaviors. Here we use the control on length-functions from §6 and will need infinite hierarchy paths with narrow ending data and a prescribed list of large domains, these hierarchy paths are provided in the appendix. Our examples will be obtained as limits of WP geodesic segments with ending data on a hierarchy path with a prescribed list of large domains.

In section 8 we use ruled surface arguments to prove an ending lamination theorem for WP geodesic rays with narrow ending data and bounded annular coefficients. A new ingredient in this section is a geometric limit argument.

The appendix, section 9, is allocated to construction of hierarchy paths with a prescribed list of large domains.

Notation 1.6. We introduce the following notations which we will use through this paper:

We denote a constant depending on other constants as a function of them. For example, $s(D, R)$ means that s is a constant depending on constants D and R.

Given $C \geq 0$ and $K \geq 1$, for functions f and g by $f \asymp_{K,C} g$ we mean that

$$\frac{1}{K} g(x) - C \leq f(x) \leq K g(x) + C$$

for every x in the domain of the functions f and g holds.

Acknowledgment: I am so grateful to my thesis advisor Yair Minsky for so many invaluable discussions through which this work has evolved. I would like to thank Jeffery Brock who generously contributed some of the ideas in section 4. I would like to thank Scott Wolpert for several communications about the behavior of Weil-Petersson geodesics near completion strata.

2 Curve complexes and hierarchy machine

Curve complex of a surface: Let $S = S_{g,b}$ be an orientable, compact surface with genus g and b boundary components, we define $\xi(S) = 3g - 3 + b \geq 1$ to be the complexity of the surface.

The curve complex $\mathcal{C}(S)$ serves to organize the isotopy classes of essential, non-peripheral, simple closed curves on S.

We start with a surface with complexity at least 1, excluding annuli and three holed spheres, to each isotopy class of essential, non-peripheral simple closed curves on S is associated a vertex ($0-$simplex) in $\mathcal{C}(S)$ and a $k-$simplex is associated to $k+1$ isotopy classes which can be realized disjointly on S. We denote the $k-$skeleton of this complex by $\mathcal{C}_k(S)$.

There are two exceptional cases, one holed torus and four holed sphere, where $1-$simplices (edges) correspond to curves intersecting respectively once and twice, which is the minimum possible intersection for curves on there surfaces.

We make the complex a metric space be making each simplex Euclidean with side lengths 1.

By the main result of [MM99] $\mathcal{C}(S)$ equipped with $d_{\mathcal{C}(S)}$ is a $\delta-$hyperbolic space in the sense of Gromov, where δ depends only on topological type of S.

Notation 2.1. We introduce the following notations and conventions about isotopy classes of curves and subsurfaces of S:

A curve on S is the isotopy class of a non-peripheral, essential simple closed curve on S, that is a vertex of $\mathcal{C}$(S). A multi-curve is the vertices of a simplex in $\mathcal{C}(S)$, that is a collection of pair-wise disjoint curves. A subsurface of S is the isotopy class of a subsurfaces of S.

Two isotopy classes of curves α and β are disjoint if they can be realized disjointly on S. Otherwise, they *overlap* each other which is denoted by $\alpha \pitchfork \beta$. Two multi-curves σ and σ' are disjoint if any pair of curves $\alpha \in \sigma$ and $\alpha' \in \sigma'$ be disjoint, otherwise σ and σ' overlap.

Laminations: A closed subset of a hyperbolic surface foliated by geodesics is a *geodesic lamination.* We denote the space of geodesic laminations of S by $\mathcal{GL}(S)$.

Each geodesic lamination can be equipped with a measure on transversal arcs invariant under isotopies of the surface relative to the lamination. We denote the space of measured geodesic laminations of S equipped with the weak* topology by $\mathcal{ML}(S)$.

$\mathcal{PML}(S)$ is the space of projective measured laminations it can be equipped with the quotient of the weak* topology on $\mathcal{ML}(S)$, for more details see for example [CEG87].

The Gromov boundary of every $\delta-$hyperbolic space has a standard topology. Klarriech shows that the there is a homeomorphism Φ from the Gromov boundary of $\mathcal{C}(S)$ to $\mathcal{EL}(S)$ the space of all *minimal filling geodesic laminations* of S and is equipped with the topology it inherits as a subset of $\mathcal{PML}(S)$ with the above topology.

Here we elaborate a result of Klarriech which describes the relation between a sequence of curves going off to the Gromov boundary of $\mathcal{C}(S)$ and the accumulation points of the sequence in $\mathcal{PML}(S)$.

Theorem 2.2. *([Kla] Theorem 1.4) If a sequence of vertices in $\mathcal{C}_0(S)$ limits to a point x in the Gromov boundary of $\mathcal{C}(S)$ then regarding each element of the sequence as a projective measured lamination every accumulation points of the sequence in $\mathcal{PML}(S)$ is topologically equivalent to $\Phi(x)$. That is their supporting geodesic laminations are topologically equivalent.*

An *annular subsurface* is an annulus Y with incompressible core in S, which is not isotopic into a boundary of S. The purpose of defining complexes for such annuli is to keep track of Dehn twisting around their cores, hence one would like $\mathcal{C}(Y)$ to be $\mathbb{Z}$. However there seems to be no natural way to do this, and we will be content with something more complicated which is nevertheless quasi-isometric to $\mathbb{Z}$.

Let $\widetilde{Y}$ be the annular cover of S to which Y lifts homeomorphically. There is a natural compactification of $\widetilde{Y}$ to a closed annulus $\widehat{Y}$ obtained in the usual way from the compactification of the universal cover $\widetilde{S} = \mathbb{H}^2$ by the closed disk. Define the vertices of $\mathcal{C}(Y)$ to be the paths connecting the two boundary components of $\widehat{Y}$, modulo isotopies that fix the endpoints (isotopy classes of arcs relative to boundary). We put an edge between any two vertices which have representatives with disjoint interiors. As before we can make $\mathcal{C}(Y)$ into a metric space with edge lengths 1.

Let $\alpha \in \mathcal{C}_0(S)$ be the core curve of Y, we write $Y = A(\alpha)$, $\mathcal{C}(\alpha) = \mathcal{C}(Y)$.

The following notions of subsurface projection and subsurface coefficient are basic in the Masur-Minsky machinery of curve complexes and hierarchical structures on the pants and marking graphs.

Subsurface projection: Again we treat annular and non-annular subsurfaces separately. Given a non-annular subsurface $Y \subseteq S$ we define

$$\pi_Y : \mathcal{GL}(S) \to \mathcal{P}(\mathcal{C}_0(Y))$$

where $\mathcal{P}(\mathcal{C}_0(Y))$ stands for the power set of $\mathcal{C}_0(S)$ as follows: Given $\lambda \in \mathcal{GL}(S)$, let $\lambda \cap Y$ be the collection of all essential curves and arcs obtained by intersecting λ and Y. Then identify all the arcs isotope to each other relative to the boundary of Y, also curves isotope to each other. The the projection would be the boundary of any regular neighborhood of this collection of curves and arcs union ∂Y.

Note that since $\mathcal{C}_0(S) \subset \mathcal{GL}(S)$ then in particular we have a projection

$$\pi_Y : \mathcal{C}_0(S) \to \mathcal{P}(\mathcal{C}_0(Y))$$

Now we continue by annular subsurfaces. We would define the projection as follows:

If γ is a simple closed curve in S crossing the core of Y transversely, then the lift of γ to $\widehat{Y}$ (the compactification of the annular cover of Y denoted by $\tilde{Y}$) has at least one component that connects the two boundaries of $\widehat{Y}$. Then together these components make up a (finite) set of diameter 1 in $\mathcal{C}(Y)$. Let $\pi_Y(\gamma)$ be this set. If γ does not intersect Y essentially (including the case that γ is the core of Y) then $\pi_Y(\gamma) = \emptyset$, as in the case of non-annular subsurfaces.

Subsurface coefficient: Now given $\sigma, \sigma' \subset \mathcal{C}(S)$ and $Y \subseteq S$ we consider the following notion of distance

$$d_Y(\sigma, \sigma') = \min\{d_{\mathcal{C}(Y)}(\pi_Y(\gamma), \pi_Y(\gamma')) : \gamma \in \sigma, \gamma' \in \sigma'\} \tag{2.1}$$

which provides a useful notion of complexity for curves systems. We it the Y subsurface coefficient of σ and σ'.

For an annular subsurface Y with core curve α we denote $\pi_Y = \pi_\alpha$ and define the annular coefficients of $\sigma, \sigma' \subset \mathcal{C}(S)$ as in (2.1), moreover we would denote it by $d_\alpha(\sigma, \sigma')$.

Notation 2.3. We introduce the following notations to related (isotopy classes of) subsurfaces of a surface S:

$Y \subseteq Z$ simply means Y is a subsurface of Z.

Given a multi-curve $\sigma \pitchfork Y$ means that σ can not be isotoped to the complement of Y. $Y \pitchfork Z$ means that $\partial Y \pitchfork Z$ and $\partial Z \pitchfork Y$, in this case we say Y and Z overlap.

2.1 The pants and marking graph

A *pants decomposition* P on a surface S is a maximal collection of disjoint curves in $\mathcal{C}(S)$. A *(partial) marking* μ consists of a pants decomposition which is the base of the marking and is denoted by $\mathrm{base}(\mu)$ together with an element of $\mathcal{C}(\alpha)$ for (some) every $\alpha \in \mathrm{base}\,\mu$. Any such element is represented a curve β called by the transversal curve to α. β intersects no other curve in $P - \alpha$ and has a minimal intersection number 1 or 2 with α depending on $S - (P - \alpha)$ be respectively a once punctured torus or four holed sphere.

By a result of Bres, see [Bus10], given any S there is a constant L_S (Bers constant) such that each $x \in \mathrm{Teich}(S)$ posses a pants decomposition (Besr pants decomposition) such that any curve in that pants decomposition has a geodesic representative of length at most L_S. Moreover, any curve in a Bers pants decomposition id called a Bers curve. Note that the constant only depends on the topological type of S. By the Collar Lemma there are only finitely many Bers pants decompositions on the surface S.

A *Bers marking* is obtained from a Bers pants decomposition by adding transversal curves of minimal length.

Pants graph: It is a graph $P(S)$ each vertex of which is the isotopy class of a pants decompositions, each edge correspond to two pants decompositions which differ by an elementary move. Pants decomposition P and P' differ by an elementary move if P' is obtained from P by fixing all the curves in P and replacing one curve α with another curve α' which have minimal intersection number (1 or 2). Giving each edge length 1 defines a distance denoted by d on $P(S)$. It is almost immediate from the definition of elementary moves on pants decompositions that for any $P, Q \in P(S)$ and any non-annular subsurface $Y \subseteq S$

$$d_Y(P,Q) \leq d(P,Q) \tag{2.2}$$

We will use this inequality frequently later on in this paper.

Marking graph: It is the graph $M(S)$ whose vertices correspond to markings and its edges correspond to markings which differ by an elementary move. An elementary move on marking roughly speaking is either an elementary move on the base of marking or is an interchange of a curve in the base and its transversal. Giving length one to each edge makes marking graph a metric space.

The following theorem plays an important role in organizing the so called tight geodesics in curve complexes of a surface and its subsurfaces and inductive construction of hierarchies in the pants and marking graphs of a surface.

Theorem 2.4. *(Bounded geodesic image)[MM00]*

There is a constant G depending on topological type of S with the following property, given $Y \subsetneq S$ and a geodesic $g : J \to \mathcal{C}(S)$ such that for every $j \in J$ $\pi_Y(g(j)) \neq \emptyset$ ($g(j) \pitchfork Y$) then

$$\mathrm{diam}_{\mathcal{C}(Y)}(\{g(j)\}_{j \in J}) \leq G$$

In the rest of this subsection we recall some results in the context of pants and marking graphs which we use often in this paper.

Distance formula: There is $M_1 > 0$ such that for any $A > M_1$, there are constants $K > 1$ and $C > 0$ such that:

$$d(P,Q) \asymp_{K,C} \sum_{\substack{Y \subseteq S \\ \text{non-annular}}} \{d_Y(P,Q)\}_A \tag{2.3}$$

where $d(P,Q)$ denotes the pants distance between $P, Q \in P(S)$ and the sum is over all non-annular subsurfaces. Here we denote
$\{a\}_A = \left\{ \begin{array}{ll} a & A \leq a \\ 0 & A > a \end{array} \right.$

Remark 2.5. Distance formula (2.3) holds for the marking distance as well summing up over all subsurfaces, including annular ones.

Theorem 2.6. *(Behrstock inequality)[Beh06]*
There is a positive constant B_0 with the property that for $W, Z \subseteq S$ such that $W \pitchfork Z$ and a partial marking μ such that $\mu \pitchfork U, V$ we have

$$\min\{d_Y(\partial Z, \mu), d_Z(\partial Y, \mu)\} \leq B_0$$

We recall the consistency theorem from [BKMM12]. Note that there this theorem is stated and proved for markings. It is straightforward to see that all the arguments go through for pants decompositions by considering only non-annular subsurfaces (excluding all annular subsurfaces).

Theorem 2.7. *(Consistency)([BKMM12] Theorem 4.3)*
Given $F_1, F_2 \geq 1$, there is F such that if a tuple $(x_Y)_{Y \subseteq S}$ of $x_Y \in \mathcal{C}(Y)$ where Y is non-annular satisfy the following two conditions:

1. *If $Y \pitchfork Z$ then $\min\{d_Y(x_Y, \partial Z), d_Z(x_Z, \partial Y)\} \leq F_1$, and*
2. *If $Y \subseteq Z$ then $d_Y(x_Y, \pi_Y(x_Z)) \leq F_2$*

Then there is $P \in P(S)$ such that for every $Y \subseteq S$

$$d_Y(P, x_Y) \leq F$$

Remark 2.8. Note that given a partial marking μ, the tuple of $x_Y = \pi_Y(\mu)$ satisfies the above two conditions with $F_1 = B_0$ and $F_2 = 1$, where B_0 is the Behrstock constant.

Definition 2.9. (Partial order in pants and marking graphs) Fix a tuple $(x_Y)_{Y \subseteq S}$ satisfying the consistency conditions in Theorem 2.7. Then we define the following two notions of partial orders with respect to $(x_Y)_{Y \subseteq S}$:

(i) By $Y \prec_k Z$, we mean that $Y \pitchfork Z$ and $d_Y(x_Y, \partial Z) \geq k(F_1 + 4)$.

(ii) By $Y \ll_k Z$, we mean that $Y \pitchfork \partial Z$ and $d_Y(x_Y, \partial Z) \geq k(F_1 + 4)$

It is almost immediate from the definition that $Y \prec_k Z \Longrightarrow Y \ll_k Z$, but not the other way round.

Also if $k > l$ then $Y \prec_k Z \Longrightarrow Y \prec_l Z$, and similarly $Y \ll_k Z \Longrightarrow Y \ll_l Z$.

As we will see in Proposition 2.14 these notions of partial order are closely related to a partial *time order* along hierarchy paths. The following theorem provides for some useful transitivity properties of the partial order defined above.

Theorem 2.10. *([BKMM12] Lemma 4.1)*

Given an integer $k > 1$, let $(x_Y)_{Y \subseteq S}$ be a tuple satisfying the consistency conditions and consider the partial order defined by it.

Let U, V and W be subsurfaces such that $x_U, x_V, x_W \neq \emptyset$ then we have:

1. *If $U \prec_k V$ and $V \ll_2 W$ then $U \prec_{k-1} W$, also if μ is a partial marking, and if $U \prec_k V$ and $V \ll_2 \mu$, then $U \prec_{k-1} \mu$.*

2. *If $U \ll_k V$ and $V \ll_2 W$ then $U \ll_{k-1} W$, also if μ is a partial marking, and if $U \ll_k V$ and $V \ll_2 \mu$ then $U \ll_{k-1} \mu$.*

3. *If $U \pitchfork V$ and both $U \ll_k \mu$ and $V \ll_k \mu$ for a partial marking μ, then U and V are $\prec_{k-1}$ −ordered, that is, either $U \prec_{k-1} V$ or $V \prec_{k-1} U$.*

2.2 Hierarchies and their resolutions in the pants and marking graph

The complete hierarchies of *tight geodesics* in curve complexes of subsurfaces of a surface were introduce by Masur and Minsky (see [MM00], [Min10], [BKMM12]). A hierarchy is the result of an inductive construction by considering geodesics in the curve complex of all subsurfaces showing up as the complement of curves along a geodesic in the curve complex of a surface. Although in most of this paper we use only hierarchy resolution paths, having a good picture of the structure of hierarchies themselves would be extremely useful for the reader to follow our arguments in this paper.

Given a geodesic h in $\mathcal{C}(Y)$, some times we refer to Y as the support of h and denote it by $D(h)$. Let (ν^-, ν^+) be a pair of partial making or lamination in S, a *complete Hierarchy of tight geodesics H* with ending data (ν^-, ν^+) is a collections of tight geodesics in the curve complex of S and the curve complexes of its subsurfaces, in what follows we list the properties which describe the inductive and layered structure of the hierarchy H, see the above references for more details:

1. There is a unique main geodesic g_S with $D(g_S) = S$, whose endpoints lie on base(ν^-) and base(ν^+).

2. For any geodesic $h \in H$ other than g_S, there exists another geodesic $k \in H$ such that, for some simplex $v \in k$, $D(h)$ is either a component of $D(k) - v$, or an annulus whose core is a component of v. We say that $D(h)$ is a *component domain* of k.

3. A subsurface in S can occur as the domain of at most one geodesic in H.

Infinite hierarchies: An infinite hierarchy is one in which there are infinite rays or lines in the curve complexes of subsurfaces. Note that in this case each of $\nu^{\pm}$ can the union of minimal, filling laminations supported on disjoint subsurfaces. Then each lamination is the limit of an infinite geodesic in the curve complex of a subsurface.

The so called *hierarchy resolution paths* in pants and marking graphs would comprise a set of transitive quasi-geodesics in the pants (marking) graphs of the surfaces with constants depending only on the topological type of S. In §5 we will characterize all stable hierarchy resolution paths in the pants graph.

A pants graph resolution of a hierarchy H is a path in the pants graph denoted by $\rho : [n, m] \to P(S)$, where $[m, n] \subset \mathbb{Z}$. It has the property that for any $i \in [m, n]$, $\rho(i)$ and $\rho(i{+}1)$ differ by an elementary move (have pants distance 1). Similarly a marking graph resolution of H is a path of clean markings such that any two consecutive markings differ by an elementary move. For a hierarchy path we denote $|\rho| = \{\rho(i) : i \in [n, m]\}$ which is a subset of $P(S)$.

Any hierarchy path consists of *slices* of a hierarchy. Each slice is union of vertices from geodesics whose supporting domains consist a tower of nested domains, note that domains are not necessarily connected.

In the following theorem we list properties of these quasi-geodesics, see also [BMM11], the main feature of these properties is that are encoded in the ending data and their subsurface coefficients:

Theorem 2.11. *(Properties of pants hierarchy resolution paths)*

Given (ν^-, ν^+), partial markings or laminations, there are hierarchy (resolution) paths $\rho : [m, n] \to P(S)$ $([m, n] \subseteq \mathbb{N})$ with ending data (ν^-, ν^+) satisfying the following list of properties:

1. *(Component domain) There are non-annular subsurfaces $Y \subseteq S$ and connected interval $J_Y \subset [m, n]$ such that for every $i \in J_Y$ $\partial Y \subset \rho(i)$.*

2. *There is an $M_1 > 0$ such that an essential, non-annular subsurface $Y \subseteq S$ with $d_Y(\nu^-, \nu^+) \geq M_1$ is a component domain of any ρ with ending data (ν^-, ν^+).*

3. *Given a component domain Y, there is a geodesic $g_Y \subset \mathcal{C}(Y)$ such that for each $j \in J_Y$ there is $v \in g_Y$ such that $v \in \rho(j)$ $(v \in g_Y \cap \rho(j))$.*

4. *(Monotonicity) Let $i, j \in J_Y$ and $i < j$ then $\pi_Y(\rho(i)) < \pi_Y(\rho(j))$ as vertices on g_Y.*

5. *There is $M_2 > 0$ with the following property: Given Y and W be component domains of ρ with $W \pitchfork Y$ let $J_Y = [j^-, j^+]$, then for every $i \in J_W$ either $d_Y(\rho(i), \rho(j^-)) \leq M_2$ or $d_Y(\rho(i), \rho(j^+)) \leq M_2$.*

   In particular, if $i > j^+$ then $d_Y(\rho(i), \rho(j^+)) \leq M_2$, and if $i < j^-$ then $d_Y(\rho(i), \rho(j^-)) \leq M_2$.

6. *For any subsurface $W \subseteq S$, for any $i \in [m,n]$ there is $x \in \mathrm{hull}_W(\nu^-,\nu^+)$ such that*
   $$d_W(\rho(i),x) \le M_2$$
   also for any $x \in \mathrm{hull}_W(\nu^-,\nu^+)$ there is $i \in [m,n]$ such that the above bound holds. Here $\mathrm{hull}_W(\nu^-,\nu^+)$ is the convex hull of $\pi_W(\nu^-)$ and $\pi_W(\nu^+)$ in $\mathcal{C}(W)$.

   In other words, $\mathrm{hull}_W(\nu^-,\nu^+)$ and $\pi_W(|\rho|)$ have M_2 Hausdorff distance in $\mathcal{C}(W)$.

Properties (1), (3) and (4) in the list follow from the structure of hierarchies and how we resolve them. Properties (2) and (5) follow from Lemma 6.1 (large link) in [MM00], Property (6) is established in the proof of Lemma 5.14 in [Min10].

Remark 2.12. A marking hierarchy resolution path $\bar{\rho} : [m,n] \to M(S)$ ($[m,n] \subseteq \mathbb{N}$) with ending data (ν^-,ν^+) satisfies the same list of properties as pants hierarchy resolution paths except in properties (1)-(6), subsurfaces Y and W could be an annular domain as well.

Definition 2.13. (Time order) Given component domains Y and W of a hierarchy path ρ we say $Y < W$ if

$$\sup J_Y < \sup J_W$$

The above notion of partial time order of component domains of a hierarchy path is closely related to the partial orders introduced in Definition 2.9, more precisely we have the following proposition:

Proposition 2.14. *Let $M = \max\{M_1, M_2, B_0\}$. Given partial markings μ and μ'. If $d_Y(\mu,\mu') > M_1$, $d_Y(\mu,\mu') > M_1$ and $Y \pitchfork W$ then either*

1. *$Y \prec_M W$ with respect to μ and $W \prec_M Y$ with respect to μ' and $Y < W$, or*

2. *$W \prec_M Y$ with respect to μ and $Y \prec_M W$ with respect to μ and $W < Y$*

This proposition follows from Lemma 4.18 in [MM00].

2.3 $\Sigma-$hulls and their projections

In this section we recall the notion of $\Sigma-$hull and projection on it from [BKMM12]. This projection is coarsely the closest point projection on the $\Sigma-$hull as a subset of the pants graph of the surface. Note that in [BKMM12] theses notions are introduced in the context of marking graphs. Other places were variations of this projection have been used are [Beh06],[BM08],[BMM11].

Given pair of partial markings or laminations (ν^-,ν^+) and $\epsilon > 0$ define

- $\Sigma_\epsilon(\nu^-,\nu^+) = \{P \in P(S) : d_W(P, \mathrm{hull}_W(\nu^-,\nu^+)) \le \epsilon$ for every non-annular $W \subseteq S\}$

where $\operatorname{hull}_W(\nu^-,\nu^+)$ is the set of all geodesics in $\mathcal{C}(W)$ connecting $\pi_W(\nu^-)$ to $\pi_W(\nu^+)$. Note that because of hyperbolicity of $\mathcal{C}(W)$ all of the geodesics connecting $\pi_W(\nu^-)$ and $\pi_W(\nu^+)$ fellow-travel each other in $\mathcal{C}(W)$.

Theorem 2.15. *([BKMM12] Proposition 5.2) There is F depending only on the topological type of the surface such that for every $\epsilon > F$ there is a projection*

$$\Pi : P(S) \to \Sigma_\epsilon(\nu^-,\nu^+)$$

with the following properties:

1. *For every non-annular subsurface $Y \subseteq S$*

$$d_Y(\Pi P, \operatorname{hull}_W(\nu^-,\nu^+)) \leq F$$

2. $\Pi\big|_{\Sigma_\epsilon(\nu^-,\nu^+)}$ *is uniformly close to the identity.*
3. Π *is coarse-Lipschitz.*

Remark 2.16. This theorem has been stated in [BKMM12] in a slightly different way also it has been stated and proved in the context of marking graph. But again it is straightforward to see that all of their arguments go through in the context of pants graph excluding all annular subsurfaces.

They show that given ending data (ν^-,ν^+) there are constants F_1 and F_2 such that the tuple $(x_Y)_{Y\subseteq S}$ where each $x_Y \in \mathcal{C}(Y)$ is a nearest point to $\pi_Y(P)$ on $\operatorname{hull}_Y(\nu^-,\nu^+)$ satisfies the consistency conditions in Theorem 2.7. Therefore there is a constant F and a pants decomposition denoted by ΠP satisfying (1).

Remark 2.17. Using the distance formula (2.3) ΠP coarsely minimizes the distance between P and the $\Sigma_\epsilon(\nu^-,\nu^+)$ in the pants graph.

3 The Weil-Petersson metric and its synthetic properties

The *Teichmüller space* of a surface S consists of all complete, finite area marked hyperbolic metrics on the surface up to isotopy. A marking of a hyperbolic surface x is a homeomorphism $(h : S \to x)$, then two marked hyperbolic surfaces $(h_1 : S \to x_1)$ and $(h_2 : S \to x_2)$ are isotopy equivalent if there is an isometry $f : x_1 \to x_2$ such that $f \circ h_1$ is isotopic to h_2.

We denote the Teichmüller space of S by $\operatorname{Teich}(S)$.

The *mapping class group* of the surface S, which we denote it by $\operatorname{Mod}(S)$ consists of all orientation preserving diffeomorphisms of S up to isotopy,

$$\operatorname{Mod}(S) = \operatorname{Diff}^+(S)/\operatorname{Diff}^0(S)$$

$\operatorname{Mod}(S)$ acts on $\operatorname{Teich}(S)$ by remarking as follows:

An element $f \in \operatorname{Mod}(S)$ maps the marked hyperbolic surface $(h : S \to x)$ to the marked hyperbolic surface $(f^{-1} \circ h : S \to x)$.

The *moduli space* of S is the space of hyperbolic metrics on S up to isotopy

$$\mathcal{M}(S) = \operatorname{Teich}(S)/\operatorname{Mod}(S)$$

Given $\epsilon > 0$ the $\epsilon-$thick part of Teichmüller space

$$\{x \in \operatorname{Teich}(S) : \operatorname{inj}(x) \geq \epsilon\}$$

where $\operatorname{inj}(x)$ is the injectivity radius of the hyperbolic surface x. The $\epsilon-$thin part is $\{x \in \operatorname{Teich}(S) : \operatorname{inj}(x) \leq \epsilon\}$.

Moreover, given a multi-curve σ (a simplex in $\mathcal{C}(S)$) we define the following regions in Teichmüller space:

- $V_\epsilon(\sigma) = \{x \in \operatorname{Teich}(S) : \ell_\alpha(x) \leq \epsilon$ for any $\alpha \in \sigma\}$
- $U_{\epsilon,\epsilon'}(\sigma) = \{x \in \operatorname{Teich}(S) : \ell_\alpha(x) \leq \epsilon$ for any $\alpha \in \sigma$ and $\ell_{\alpha'}(x) > \epsilon'$ for every $\alpha' \notin \sigma\}$

Here we recall some properties of the Weil-Petersson metric and its geodesics which will be used in this paper. Some reference for these material are [Wol10], [Wol08], [Wol03], [McM00], [DW03].

Given $\varphi, \psi \in \mathcal{QD}(x) = T^* \operatorname{Teich}(S)$ the Weil-Petersson L^2 co-product is defined by

$$\mathcal{R}e\Big(\int_S \varphi\overline{\psi}\rho^{-2}\Big)$$

where $\rho(z)^2|dz|^2$ is the hyperbolic metric of the marked hyperbolic surface $(h : S \to x)$.

This co-product induces a norm on Teichmüller space via the standard pairing of quadratic differentials and Beltrami differentials defined by $\int_S \varphi\mu$.

The Weil-Petersson metric is defined by the polarization of this norm on Teichmüller space. In this paper we are interested in study of the global behavior of geodesics of this metric.

The Weil-Petersson metric is a Riemannian metric with negative sectional curvatures which is invariant under the action of the mapping class group of the surface.

It is an incomplete metric, but still geodesically convex which together with the negative curvature implies that the completion of Teichmüller space with respect to it, denoted by $\overline{\operatorname{Teich}(S)}$, is a CAT(0) space, for background about CAT(0) space see for example [BH99].

Length-functions: Given any $\alpha \in \mathcal{C}(S)$ the $\alpha-$length-function

$$\ell_\alpha : \operatorname{Teich}(S) \to \mathbb{R}^+$$

assigns to $x \in \operatorname{Teich}(S)$ the length of the geodesic representative of α on the marked hyperbolic surface x.

There is an extension of the notion of length-function to measured lamination, see [CEG87], for a measured lamination μ we denote the $\mu-$length-function by $\ell_\mu(x)$.

Notation 3.1. For a multi-curve σ, a simplex in $\mathcal{C}_0(S)$, and $x \in \text{Teich}(S)$ we define

$$\ell_\sigma(x) = \max_{\alpha\in\sigma} \ell_\alpha(x)$$

Fenchel-Nielsen coordinates and twist parameters: Given a pants decomposition $(\ell_\gamma, \theta_\gamma)_{\gamma\in P}$ is the standard Fenchel-Nielsen coordinate which maps $\text{Teich}(S)$ to $\prod_{\gamma\in P}\mathbb{R}^+_\gamma \times \mathbb{R}_\gamma$. The first coordinate of the pair $\mathbb{R}^+_\gamma \times \mathbb{R}_\gamma$ is the $\gamma-$length-function and the second coordinate is the twist parameter of γ. For more details about Fenchel-Nielsen coordinates and twit parameters see [Bus10] §3.

The Weil-Petersson completion of Teichmüller space and the completion strata: The incompleteness of the Weil-Petersson metric corresponds to finite-length paths in Teichmüller space along which length-functions of curves converge to zero [Wol10]. In [Mas76], Masur gives a concrete description of the completion as the *augmented Teichmüller space*, obtained from Teichmüller space by adding strata consisting of spaces $\mathcal{S}(\sigma)$ defined by the vanishing of length-functions $\{\ell_\alpha = 0\}$, for each $\alpha \in \sigma$ where σ is a simplex in the augmented curve complex $\widehat{\mathcal{C}(S)} = \mathcal{C}(S) \cup \emptyset$. Points in the $\sigma-$stratum, $\mathcal{S}(\sigma)$, correspond to nodal Riemann surfaces z, where (paired) cusps are introduced along the curves in σ.

One can describe the topology via extended Fenchel-Nielsen coordinates as follows: Given a pants decomposition P we extend the Fenchel-Nielsen coordinates $(\ell_\gamma, \theta_\gamma)_{\gamma\in P}$ as above to allow length-function take value 0 as well. Then take the quotient by identifying $(0, \theta) \sim (0, \theta')$ in each $\mathbb{R}^+ \times \mathbb{R}$ factor. The topology near any point of a stratum $\mathcal{S}(\sigma)$, where $\sigma \subset P$, is such that this map is a homeomorphism near that point.

Then the stratum $\mathcal{S}(\sigma)$ is naturally the product of lower dimensional Teichmüller spaces corresponding to the complete, finite area hyperbolic pieces of the nodal surface $z \in \mathcal{S}(\sigma)$.

Theorem 3.2. *(Non-refraction)[DW03],[Wol03], [Yam04]*

Let $\zeta : [0, T] \to \overline{\text{Teich}(S)}$ be a WP geodesic segment, and let σ^- and σ^+ be the maximal simplicies in $\mathcal{C}(S)$ such that $\zeta(0) \in \mathcal{S}(\sigma^-)$ and $\zeta(T) \in \mathcal{S}(\sigma^+)$ then

$$int(\zeta) \subset \mathcal{S}(\sigma^- \cap \sigma^+)$$

As a consequence of non-refraction theorem Daskalopoulos and Wentworth in [DW03] and Wolpert in [Wol03] show that any pseudo-Anosov element of the mapping class group f has an axis in Teichmüller space equipped with the WP metric i.e. there is a bi-infinite geodesic $Ax_f \subset \text{Teich}(S)$ such that

$$d_{\text{WP}}(x, fx) = \inf_{y\in\text{Teich}(S)} d_{\text{WP}}(y, fy)$$

for every $x \in \mathrm{Ax}_f$.

Through the following theorem of Brock the hierarchies of curves complexes and their resolution in pants and marking graphs play the main role in our study of the global behavior of WP geodesics:

Theorem 3.3. *(Quasi-isometric model)[Bro03]*

There are constants $K_{\mathrm{WP}} > 1$ and $C_{\mathrm{WP}} > 0$ depending on topological type of S, such that the map

$$Q : \overline{\mathrm{Teich}(S)} \to P(S)$$

which assigns to a marked hyperbolic surface x a Bers pants decomposition on x is a $(K_{\mathrm{WP}}, C_{\mathrm{WP}})-$quasi-isometry.

Notation 3.4. In this paper, given $x \in \overline{\mathrm{Teich}(S)}$, we denote by $Q(x)$ a Bers pants decomposition of x and by $\mu(x)$ a Bers marking of x. The base of a Bers marking is a Bers pants decomposition and the transversal curve to each curve in the base has minimum length among all transversals to that curve.

Gradient of length-functions: Using Riera's formula for the WP paring of quadratic differentials Wolpert gives the following estimate for pairing of gradient of length-functions:

Lemma 3.5. *[Wol08] The WP pairing of length-function gradients of disjoint geodesics α, β satisfies*

$$0 < \langle \mathrm{grad}\, \ell_\alpha, \mathrm{grad}\, \ell_\beta \rangle - \tfrac{2}{\pi} \ell_\alpha \delta_{\alpha\beta} = O(\ell_\alpha^2 \ell_\beta^2)$$

where for c_0 positive the remainder term constant is uniform for $\ell_\alpha, \ell_\beta \leq c_0$.

Corollary 3.6. *Given l and a positive with $l > a$, there is $\boldsymbol{r} = \boldsymbol{r}(l,a)$ such that if $\ell_\alpha(x) \leq l - a$ and $\ell_\alpha(x') \geq l$ then $d_{\mathrm{WP}}(x', x) \geq \boldsymbol{r}$.*

Proof. By the above lemma if $\ell_\alpha(p) \leq c_0$ then

$$|| \mathrm{grad}\, \ell_\alpha(p) ||_{\mathrm{WP}} \leq (\frac{2}{\pi} c_0 + C(c_0) c_0^4)^{1/2}$$

Let u be the WP geodesic segment parametrized by arc-length from x to x', let t^* be the first time that $\ell_\alpha(u(t)) = l$, then $\ell_\alpha(u(t)) \leq l$ for every $t \in [0, t^*]$, thus we have

$$a \leq \textstyle\int_0^{t^*} || \mathrm{grad}\, \ell_\alpha(u(t)) ||_{\mathrm{WP}} dt \leq (\tfrac{2}{\pi} l + C(l)(l)^4)^{1/2} t^*$$

so $t^* \geq \frac{a}{\left(\frac{2}{\pi} l + C(l) l^4\right)^{1/2}}$ and therefore for $\mathbf{r}(l,a) = \frac{a}{\left(\frac{2}{\pi} l + C(l) l^4\right)^{1/2}}$ the corollary follows. ☐

Using the estimates on pairings of the gradient of length-functions Wolpert gives an asymptotic expansion for the WP metric near completion strata. These expansion shows that there are asymptotic quasi-flats transverse to strata corresponding to any pair of pinching curves.

Tangent cones at the Weil-Petersson completion strata: The completion of Teichmüller space with the Weil-Petersson metric is a CAT(0) space. At any point in a CAT(0) space one can consider the Alexandrov tangent cone consisting of equivalence classes of geodesics starting at that point which have 0 initial angle in the sense of Alexandrov, for more detail see [BH99].

Given σ a simplex in $\mathcal{C}(S)$, let χ be full marking on $S-\sigma$, consider the map

$$\mathcal{L}(\zeta(t)) = (\ell_\alpha^{1/2}(\zeta(t)), \ell_\beta^{1/2}(\zeta(t)))_{\alpha\in\sigma,\beta\in\chi}$$

where $\zeta : [0,T] \to \overline{\mathrm{Teich}(S)}$ is a geodesic segment with $\zeta(0) = p \in \mathcal{S}(\sigma)$. Then define

$$\Lambda : \zeta \to (2\pi)^{1/2}\frac{d}{dt}\Big|_{t=0}\mathcal{L}(\zeta(t))$$

Wolpert gives the following description of the Alexandrov tangent cone at p.

Theorem 3.7. *[Wol08] The map Λ from the tangent cone of the WP metric at p to $\mathbb{R}_{\geq 0}^{|\sigma|} \times T_p\,\mathrm{Teich}(S-\sigma)$ is an isometry of tangent cones with restriction of inner products. A WP terminating geodesic ζ with root length-function initial derivative $\frac{d}{dt}\big|_{t=0}\ell_\alpha^{1/2}(\zeta(t))$ vanishing is contained in the stratum $\{\ell_\alpha = 0\}$, $\mathcal{S}(\sigma) \subset \{\ell_\alpha = 0\}$.*

3.1 Ending data

In this subsection we recall the notion of ending data for WP geodesic rays introduced by Brock, Masur and Minsky in [BMM11]. We start with some background.

Theorem 3.8. *(Convexity of length-functions)[Wol08] Given $\epsilon > 0$ there is $c(\epsilon)$ with the following property: Let $g : (a,b) \to \mathrm{Teich}(S)$ be a WP geodesic parametrized by arc-length and $\alpha \in \mathcal{C}_0(S)$. If for some $t \in (a,b)$ $\mathrm{inj}\, g(t) \geq \epsilon$ ($g(t)$ be a point in the $\epsilon-$thick part of Teichmüller space) then we have*

$$\ddot{\ell}_\alpha(g(t)) \geq c(\epsilon)\ell_\alpha(g(t)) \tag{3.1}$$

Similar inequality holds for length of any measured lamination μ, that is

$$\ddot{\ell}_\mu(g(t)) \geq c(\epsilon)\ell_\mu(g(t))$$

Remark 3.9. The above estimates are local and only depend on the length of the curve α or the measured lamination μ and the injectivity radius of the surface $g(t)$.

Definition 3.10. (Ending measure) The weak* limit of any weighted sequence of distinct Bers curves along a WP geodesic ray r is an ending measure for r.

The convexity of length-functions along WP geodesics and some properties of CAT(0) spaces as was shown in [BMM11] implies that the following notion of *ending lamination* for WP geodesic rays is well-defined.

Definition 3.11. (Ending Lamination) The union of pinching curves along a WP geodesic ray and the geodesic laminations arising as supports of all ending measures of r is the ending lamination of r.

Definition 3.12. (Ending data of Weil-Petersson geodesics) For each open end of a geodesic $g : (a, b) \to \overline{\text{Teich}(S)}$ we have an ending data which is a partial marking or a lamination. We denote it by $\nu^+(g)$, the forward end, and $\nu^-(g)$, the backward end.

If g in forward end can be extended to a b such that $g(b) \in \overline{\text{Teich}}(S)$ then we let $\nu^+(g)$ denote a (partial) Bers marking of $g(b)$ (there are finitely many of them and we pick an arbitrary one). Otherwise, $\nu^+(g)$ is the ending lamination of the ray $g|_{[0,b)}$ as defined above. We define $\nu^-(g)$ similarly.

Here we recall a couple of the properties of the ending measures and laminations proved in [BMM10]:

Lemma 3.13. *(Decreasing length along WP geodesic rays) Let μ be any ending measure of a WP geodesic ray r then $\ell_\mu(r(t))$ is a decreasing function.*

Lemma 3.14. *Let $r_n \to r$ be a convergent sequence of rays in the WP visual sphere at x. Then if μ_n is any sequence of ending measures or weighted pinching curves for r_n, any representative $\mu \in \mathcal{ML}(S)$ of the limit $[\mu]$ of projective classes $[\mu_n]$ in $\mathcal{PML}(S)$ has bounded length along the ray r.*

4 Compactness arguments in the Weil-Petersson completion of Teichmüller space

In this section we study length-functions and twist parameters along sequences of bounded length WP geodesic segments in the WP completion of Teichmüller space.

In §4.2 we will prove a modified version of Lemma 4.5 in [BMM11] about the development of Dehn twists along sequences of uniformly bounded length WP geodesic segments, Theorem 4.6, further in corollaries 4.12 and 4.11 we give a somewhat quantified version of these results. These results are a kind of twist parameter versus length-function control along WP geodesic segments.

The proof uses Wolpert's characterization of limits of sequences of bounded length WP geodesic segments in the Weil-Petersson completion of Teichmüller space. In §4.1 we state Wolpert's geodesic limit theorem and using suggestions of Jeff Brock give a modification of that, Theorem 4.5. We need this modified version to prove our results in §4.2. These results play an essential role in §6 where we study the itinerary of WP geodesics fellow-traveling hierarchy paths.

4.1 Limits of sequences of uniformly bounded length WP geodesic segments

In this subsection we provide a modified version of Wolpert's geodesic limit theorem. Denote by $\mathrm{tw}(\sigma)$ the subgroup of $\mathrm{Mod}(S)$ generated by Dehn twists about the curves in a multi-curve σ. Using the non-refraction of the Weil-Petersson completion strata, Theorem 3.2, and the fact that the quotient of any $V_\epsilon(\sigma)$ by the action of $\mathrm{tw}(\sigma)$ is compact, Wolpert gives the following characterization of the limits of uniformly bounded length WP geodesic segments in the Teichmüller space.

Theorem 4.1. *[Wol03] Let $\zeta_n : [0,T] \to \mathrm{Teich}(S)$ be a sequence of WP geodesic segments of length T. Then after possibly passing to a subsequence there exists a partition of the interval $[0,T]$ by $0 = t_0 < t_1 < t_2 < ... < t_k < t_{k+1} = T$, simplices $\sigma_0, ..., \sigma_{k+1}$ and for $i = 1, ..., k+1$ simplices $\tau_i = \sigma_{i-1} \cap \sigma_i$ in $\widehat{\mathcal{C}(S)}$ such that $\tau_1 = \sigma_0$ and for each $1 \leq i < k+1$ $\tau_i \subsetneq \sigma_i$ and $\tau_{i+1} \subsetneq \sigma_i$. Moreover a piecewise geodesic*

$$\hat{\zeta} : [0,T] \to \overline{\mathrm{Teich}(S)}$$

with the following properties

1. *$\hat{\zeta}((t_{i-1}, t_i)) \subset \mathcal{S}(\tau_i)$ for $i = 1, ..., k+1$*
2. *$\hat{\zeta}(t_i) \in \mathcal{S}(\sigma_i)$, for $i = 0, ..., k+1$*
3. *There are elements $\psi_n \in \mathrm{Mod}(S)$ and $\mathcal{T}_{i,n} \in \mathrm{tw}(\sigma_i - \tau_i \cup \tau_{i+1})$, for $i = 1, .., k$ such that $\psi_n \circ \zeta_n|_{[0,t_1]}$ converges to $\hat{\zeta}|_{[0,t_1]}$ in $\overline{\mathrm{Teich}(S)}$ and similarly for each $i = 1, .., k$*

   $$\mathcal{T}_{i,n} \circ ... \circ \mathcal{T}_{1,n} \circ \psi_n \circ \zeta_n|_{[t_i, t_{i+1}]} \to \hat{\zeta}|_{[t_i, t_{i+1}]}$$

   as $n \to \infty$
4. *The elements ψ_n are either trivial or unbounded and the elements $\mathcal{T}_{i,n}$ are unbounded.*
5. *The piecewise geodesic $\hat{\zeta}$ is the minimal length path in $\overline{\mathrm{Teich}(S)}$ joining $\hat{\zeta}(0)$ to $\hat{\zeta}(T)$ and intersecting the strata $\mathcal{S}(\sigma_1), \mathcal{S}(\sigma_2), ..., \mathcal{S}(\sigma_k)$ in order.*

The following two lemmas which were suggested to us by Jeff Brock help us to considerably improve the picture of limits of uniformly bounded length WP geodesic segments described by Wolpert in Theorem 4.1, see Theorem 4.5. For similar results see [Wu].

Lemma 4.2. *Given a sequence of WP geodesic segments $\zeta_n : [0,T] \to \mathrm{Teich}(S)$, let the piecewise geodesic path $\hat{\zeta} : [0,T] \to \mathrm{Teich}(S)$, the partition $0 = t_0 < t_1 < ... < t_{k+1} = T$, simplices σ_i and τ_i be as in Theorem 4.1. Then $\tau_0 = \tau_1 = ... = \tau_{k+1}$.*

Proof. The proof is by contradiction. For $\delta < \min_{i=1,...,k+1} \frac{t_i - t_{i-1}}{2}$ by Theorem 4.1 (1) $\hat{\zeta}|_{[t_i-\delta,t_i)}$ lies in the τ_i-stratum and $\hat{\zeta}|_{(t_i,t_i+\delta]}$ lies in the $\tau_{i+1}-$stratum. Moreover, by (5) their concatenation is distance minimizing among all paths in $\overline{\text{Teich}(S)}$ connecting $\hat{\zeta}(t_i - \delta)$ to $\hat{\zeta}(t_i + \delta)$ and intersecting $\mathcal{S}(\sigma_i)$.

Now since $\hat{\zeta}|_{[t_i-\delta,t_i+\delta]}$ is distance minimizing then as Wolpert shows in [Wol08] page 328, at the point $\hat{\zeta}(t_i) \in \mathcal{S}(\sigma_i)$ we have the following equality of derivatives of the square roots of $\alpha-$length-functions for any $\alpha \in \sigma_i$

$$\frac{d}{dt}\Big|_{t=t_i^+} \ell_\alpha^{1/2}(\hat{\zeta}|_{[t_i,t_i+\delta]}) = -\frac{d}{dt}\Big|_{t=t_i^-} \ell_\alpha^{1/2}(\hat{\zeta}|_{[t_i-\delta,t_i]}) \tag{4.1}$$

Now let $\hat{\tau} = \tau_1$. If for every $i = 1, 2, ..., k+1$, $\tau_{i+1} = \tau_i$ then the lemma follows immediately.

Otherwise, let $i > 0$ be the smallest index such that $\tau_{i+1} \neq \tau_i$, thus there is a curve α such that either $\alpha \in \tau_{i+1} - \tau_i$ or $\alpha \in \tau_i - \tau_{i+1}$. In the first case since $\hat{\zeta}|_{(t_i,t_i+\delta]} \subset \mathcal{S}(\tau_{i+1})$ we have $\ell_\alpha^{1/2}(\hat{\zeta}(t)) = 0$ for all $t \in [t_i, t_i + \delta]$, thus for the curve α the right hand side of (4.1) is zero.

Let P be a pants decomposition containing σ_i (note that $\tau_i, \tau_{i+1} \subsetneq \sigma_i$). Then since $\hat{\zeta}|_{[t_i,t_i-\delta]} \subset \mathcal{S}(\tau_i)$ and $\alpha \notin \tau_{i+1}$ the description of the WP tangent cone at the completion strata as follows implies that the left hand side of (4.1) is non zero.

To see this note that by Theorem 3.7 if $\frac{d}{dt}\big|_{t=t_i} \ell_\alpha^{1/2}(\hat{\zeta}(t)) = 0$ then $\ell_\alpha^{1/2}(\hat{\zeta}(t)) = 0$ for all $t \in [t_i - \delta, t_i)$, but τ_i is the maximal simplex such that $\hat{\zeta}([t_i - \delta, t_i)) \subset \mathcal{S}(\tau_i)$ and $\alpha \notin \tau_i$ which is a contradiction.

Therefore, we get a contradiction to the equality of derivatives in (4.1).

Similarly we can get a contradiction in the case $\alpha \in \tau_i - \tau_{i+1}$. Therefore, we conclude that $\tau_i = \tau_{i+1}$ and the lemma follows.

□

In the rest of the paper we denote

$$\tau_i \equiv \hat{\tau}$$

Lemma 4.3. *Let $\zeta_n : [0, T] \to \text{Teich}(S)$ be a sequence of WP geodesic segments, let simplices σ_i for $i = 0, ..., k+1$ be as in Theorem 4.1 and the simplex $\hat{\tau}$ be as above. Then for every $\gamma_n = \varphi_{i,n}^{-1}(\gamma)$ where $\gamma \in \sigma_i - \hat{\tau}$ and $i = 1, ..., k$ the power of $\mathcal{D}_{\gamma_n}$ in $\mathcal{T}_{i,n}$ goes to ∞ as $n \to \infty$.*

Proof. For each $i = 1, ..., k$ define

$$\sigma_{i,n} = \varphi_{i,n}^{-1}(\sigma_i)$$

For each $i = 1, ..., k$ define the geodesic segments

$$\zeta_{i,n} = \varphi_{i,n} \circ \zeta_n|_{[t_{i-1},t_{i+1}]}$$

where $0 = t_1 < ... < t_i < t_{i+1} < ... < t_k$ is the partition from Theorem 4.1. We claim that

Claim 4.4. There are $\epsilon > 0$ and $\delta > 0$ such that for every n sufficiently large

(i) $\epsilon \leq \ell_{\gamma_n}(\zeta_{i,n}(t_i \pm \delta)) \leq 4\epsilon$ where $\gamma_n \in \sigma_{i,n}$, and

(ii) $\ell_\alpha(\zeta_{i,n}(t_i \pm \delta)) \geq \epsilon$ for every α disjoint from $\sigma_{i,n}$.

The claim follows from the limiting picture and continuity of length-functions.

To see this, let $\delta < \min_{i=1,\dots,k+1} \frac{t_i - t_{i-1}}{2}$. By Lemma 4.2 the points $\hat{\zeta}(t_i \pm \delta)$ are in the interior of $\mathcal{S}(\hat{\tau})$ and $\hat{\zeta}(t_i)$ is in $\mathcal{S}(\sigma_i)$. Thus the continuity of length-functions along $\hat{\zeta}$ implies that there are ϵ and δ such that:

(i') $2\epsilon \leq \ell_\gamma(\hat{\zeta}(t_i \pm \delta)) \leq 3\epsilon$ for every $\gamma \in \sigma_i - \hat{\tau}$, and

(ii') $\ell_\alpha(\hat{\zeta}(t_i \pm \delta)) > 2\epsilon$ for every $\alpha \notin \sigma_i - \hat{\tau}$

Moreover, by Theorem 4.1 (3) $\zeta_{i,n}\big|_{[t_{i-1},t_i]} \to \hat{\zeta}\big|_{[t_{i-1},t_i]}$ as parametrized geodesics. The continuity of length-functions tells us that in a converging sequence of points in the Teichmüller space equipped with the WP metric the length-functions also converge. Thus for n sufficiently large bounds (i) and (ii) at $\zeta_{i,n}(t_i - \delta)$ immediately follow from the ones in (i') and (ii').

Further, since $\varphi_{i,n} = \mathcal{T}_{i,n} \circ \varphi_{i-1,n}$ again by Theorem 4.1 $\mathcal{T}_{i,n} \circ \zeta_{i,n}\big|_{[t_i,t_{i+1}]} \to \hat{\zeta}\big|_{[t_i,t_{i+1}]}$ as parametrized geodesics. Now again by the continuity of length-functions from (i') and (ii') we get:

$\epsilon \leq \ell_{\gamma_n}(\mathcal{T}_{i,n} \circ \zeta_n(t_i + \delta)) \leq 4\epsilon$ for every $\gamma_n \in \sigma_{i,n}$ and $\ell_\alpha(\mathcal{T}_{i,n} \circ \zeta_n(t_i + \delta)) \geq \epsilon$ for every $\alpha \notin \sigma_{i,n}$.

Now $\mathcal{T}_{i,n} \in \mathrm{tw}(\sigma_{i,n} - \varphi_{i,n}^{-1}(\hat{\tau}))$ so it preserves the isotopy class and the length of every curve which does not overlap $\sigma_{i,n}$ (either is in $\sigma_{i,n}$ or is disjoint from it). Thus the bounds (i) and (ii) hold at $\zeta_{i,n}(t_i + \delta)$ as well.

This finishes the proof of our claim.

Now we continue to prove the lemma by contradiction, fix $0 \leq i \leq k+1$ and $\gamma \in \sigma_i - \hat{\tau}$. Pick β_n to be a curve on $\zeta_{i,n}(t_i - \delta)$ with minimum intersection number (1 or 2) with $\gamma_n = \varphi_{i,n}^{-1}(\gamma)$ which is disjoint from $\varphi_{i,n}^{-1}(\sigma_i - \gamma) = \sigma_{i,n} - \gamma_n$ then by the two bounds in above claim and the Collar lemma (see [Bus10] §4.1) we have

$$\ell_{\beta_n}(\zeta_{i,n}(t_i - \delta)) \leq \omega(\epsilon) + C \tag{4.2}$$

where C depends only on the topological type of S. Now for any $\gamma_n \in \sigma_{i,n}$ denote by $\mathcal{D}_{\gamma_n}$ the Dehn twist about γ_n, if the absolute value of the power of $\mathcal{D}_{\gamma_n}$ in $\mathcal{T}_{i,n}$ be bounded by N then again by the two bounds in the above claim, the definition of Dehn twist and the Collar lemma we would have

$$\ell_{\beta_n}(\zeta_{i,n}(t_i + \delta)) \leq N\epsilon + \omega(\epsilon) + C \tag{4.3}$$

Moreover, since $\ell_{\gamma_n}(\zeta_{i,n}(t_i)) \to 0$ as $n \to \infty$ and $\beta_n \pitchfork \gamma_n$ then by the Collar lemma

$$\ell_{\beta_n}(\zeta_{i,n}(t_i)) \to \infty \tag{4.4}$$

as $n \to \infty$.

But (4.4), (4.2) and (4.3) violate the convexity of the β_n-length-function along the WP geodesic segment $\zeta_{i,n}$ and we get the contradiction and the lemma follows.

$\square$

Now for the purpose of reference in this paper we state the following version of geodesic limit theorem which are the properties listed in Theorem 4.1 modified to incorporate lemmas 4.2 and 4.3.

Theorem 4.5. *(Geodesic limit) Let $\zeta_n : [0,T] \to \overline{\mathrm{Teich}(S)}$ be a sequence of WP geodesic segments of length T. Then after possibly passing to a subsequence there exists a partition of the interval $[0,T]$ by $0 = t_0 < t_1 < t_2 < ... < t_k < t_{k+1} = T$, and simplices $\sigma_0, ..., \sigma_{k+1}$ in $\mathcal{C}(S)$ such that for every i, $\sigma_i \cap \sigma_{i+1} = \hat{\tau}$ ($\sigma_0 = \hat{\tau}$) and a piecewise geodesic*

$$\hat{\zeta} : [0,T] \to \overline{\mathrm{Teich}(S)}$$

with the following properties

1. *$\hat{\zeta}((t_{i-1}, t_i)) \subset \mathcal{S}(\hat{\tau})$ for $i = 1, ..., k+1$*
2. *$\hat{\zeta}(t_i) \in \mathcal{S}(\sigma_i)$, for $i = 0, ..., k+1$*
3. *There are elements $\psi_n \in \mathrm{Mod}(S)$ and $\mathcal{T}_{i,n} \in \mathrm{tw}(\sigma_i - \hat{\tau})$, for $i = 1, .., k$ such that $\psi_n \circ \zeta_n|_{[0,t_1]}$ converges to $\hat{\zeta}|_{[0,t_1]}$ in $\overline{\mathrm{Teich}(S)}$ and for each $i = 1, .., k$,*

   $$\mathcal{T}_{i,n} \circ ... \circ \mathcal{T}_{1,n} \circ \psi_n \circ \zeta_n|_{[t_i,t_{i+1}]} \to \hat{\zeta}|_{[t_i,t_{i+1}]}$$

   as $n \to \infty$
4. *The elements ψ_n are either trivial or unbounded and for every i and $\gamma_n \in \varphi_{i,n}^{-1}(\sigma_i)$ the power of $\mathcal{D}_{\gamma_n}$ in the element $\mathcal{T}_{i,n}$ is unbounded.*
5. *The piecewise geodesic $\hat{\zeta}$ is the minimal length path in $\overline{\mathrm{Teich}(S)}$ joining $\hat{\zeta}(0)$ to $\hat{\zeta}(T)$ and intersecting the strata $\mathcal{S}(\sigma_1), \mathcal{S}(\sigma_2), ..., \mathcal{S}(\sigma_k)$ in order.*

For convenience for every $i = 0, 1, ..., k+1$ we define:

$$\varphi_{i,n} = \mathcal{T}_{i,n} \circ ... \circ \mathcal{T}_{1,n} \circ \psi_n \tag{4.5}$$

4.2 Length-function versus twist parameter control

In this subsection roughly speaking we show that the sequences of curves on surfaces along a sequence of uniformly bounded length WP geodesic segments which are getting short are the same as the sequences of curves about which a high Dehn twist develops, except possibly at the end points. This result is a modification of Lemma 4.5 in [BMM11].

Theorem 4.6. *Given positive constants* ϵ_0, T *and* s*, let* $\zeta_n : [0, T_n] \to \mathrm{Teich}(S)$ *be a sequence of Weil-Petersson geodesic segments parametrized by arc-length of length* $2s \leq T_n \leq T$*, then given a sequence of curves* α_n *we have the following:*

1. *If there are subintervals* $J_n \subseteq [s, T_n - s]$ *such that*
   - *(a)* $\sup_{t \in J_n} \ell_{\alpha_n}(\zeta_n(t)) \geq \epsilon_0$*, and*
   - *(b)* $\inf_{t \in J_n} \ell_{\alpha_n}(\zeta_n(t)) \to 0$ *as* $n \to \infty$

   then
   $$d_{\alpha_n}(\mu(\zeta_n(0)), \mu(\zeta_n(T_n))) \to \infty$$
   as $n \to \infty$*.*
2. *(a)* $\sup_{t \in [0,T_n]} \ell_{\alpha_n}(\zeta_n(t)) \geq \epsilon_0$*, and*

   (b) $d_{\alpha_n}(\mu(\zeta_n(0)), \mu(\zeta_n(T_n))) \to \infty$ *as* $n \to \infty$

   then
   $$\inf_{t \in [0,T_n]} \ell_{\alpha_n}(\zeta_n(t)) \to 0$$
   as $n \to \infty$*.*

Proof. Passing to a subsequence, trimming the intervals slightly and changing the constants we may assume that $T_n \equiv T$. Then by Theorem 4.5, after again passing to a subsequence, we have a partition of $[0, T]$ with $0 = t_0 < t_1 < ... < t_{k+1} = T$, simplices σ_i for $i = 0, 1, ..., k+1$ in $\mathcal{C}(S)$ and a simplex $\hat{\tau}$ and a piecewise geodesic path
$$\hat{\zeta} : [0, T] \to \mathrm{Teich}(S)$$
for which $\hat{\zeta}([t_i, t_{i+1}])$ is a geodesic segment in $\mathcal{S}(\hat{\tau})$ joining the stratum $\mathcal{S}(\sigma_i)$ to $\mathcal{S}(\sigma_{i+1})$ and elements $\mathcal{T}_{i,n} \in \mathrm{tw}(\sigma_i - \hat{\tau})$ are unbounded in $\mathrm{Mod}(S)$. Assume that the conclusion of Theorem 4.5 hold and let $\varphi_{i,n} \in \mathrm{Mod}(S)$ be as in (4.5).

We start by setting up some notation, for each $i = 0, ..., k+1$ and n let
$$\sigma_{i,n} = \varphi_{i,n}^{-1}(\sigma_i) = \varphi_{i+1,n}^{-1}(\sigma_i)$$
be the pull backs of σ_i to the ζ_n picture. For each $i = 1, ..., k$ and n let
$$\tau_{i,n} = \varphi_{i-1,n}^{-1}(\hat{\tau})$$
For each $i = 0, ..., k+1$, choose partial markings μ_i of $S - \sigma_i$ such that

1. $\sigma_i \subset \text{base}(\mu_i)$, and
2. μ_i restricts to a full marking of each connected component $Y \subset S - \sigma_i$ with complexity at least one.

Furthermore, for each $i = 0, ..., k+1$, let $\hat{\mu}_i$ be an enlargement of μ_i such that $\text{base}(\hat{\mu}_i) = \text{base}(\mu_i)$ and $\hat{\mu}_i$ be a full marking of each connected component of $S - \hat{\tau}$ of complexity at least 1. Note that each $\hat{\mu}_i$ differs from μ_i by addition of transversals to the components of $\sigma_i - \hat{\tau}$.

Further, for each $i = 1, ..., k+1$ define the pullbacks

$$\mu_{i,n}^{-} = \varphi_{i-1,n}^{-1}(\hat{\mu}_i)$$

and for each $i = 0, ..., k$

$$\mu_{i,n}^{+} = \varphi_{i,n}^{-1}(\hat{\mu}_i)$$

We continue to measure the twisting of these markings relative to $\gamma_n = \varphi_{i,n}^{-1}(\gamma)$ for a fixed $0 < i \leq k+1$ and $\gamma \in \sigma_i - \hat{\tau}$.

We make the following three claims:

Claim 4.7. $d_{\gamma_n}(\mu_{j,n}^{-}, \mu_{j,n}^{+})$ is bounded for every $j \neq i$.

Claim 4.8. $d_{\gamma_n}(\mu_{j,n}^{+}, \mu_{j+1,n}^{-})$ is bounded for all j.

Claim 4.9.

$$d_{\gamma_n}(\mu_{i,n}^{-}, \mu_{i,n}^{+}) \to \infty \tag{4.6}$$

as $n \to \infty$

To see Claim 4.7, note that $\text{base}(\mu_{j,n}^{+})$ and $\text{base}(\mu_{j,n}^{-})$ both contain

$$\sigma_{j,n} = \varphi_{j,n}^{-1}(\sigma_j) = \varphi_{j-1,n}^{-1}(\sigma_j)$$

Now we claim that

$$\gamma_n \notin \sigma_{j,n} \tag{4.7}$$

for any $j \neq i$

Otherwise the length of γ_n along ζ_n would converge to 0 both at t_i and at t_j, and hence by convexity on of length-functions on all of $[t_{i-1}, t_i]$ or $[t_i, t_{i+1}]$ (the first if $j < i$ and the second if $j > i$). This contradicts the choice of $\gamma \in \sigma_i - \hat{\tau}$ and Theorem 4.5 (1), so we conclude that (4.7) holds. Now γ_n intersects $\mu_{j,n}^{\pm}$ nontrivially, so $\pi_{A(\gamma_n)}(\mu_{j,n}^{\pm})$ is nonempty. Moreover $\mu_{j,n}^{+} = \mathcal{T}_{j,n}(\mu_{j,n}^{-})$ where $\mathcal{T}_{j,n} \in \text{tw}(\sigma_j)$, thus the projections of $\mu_{i,n}^{-}$ and $\mu_{i,n}^{+}$ are a bounded distance apart in $A(\gamma_n)$, establishing Claim 4.7.

To prove Claim 4.8, note that μ_j and μ_{j+1} are full markings of $S - \hat{\tau}$, where their marking distance is some finite number. Hence we may connect them with a finite sequence of full markings of $S - \hat{\tau}$. Applying $\varphi_{j,n}^{-1}$, we obtain a sequence of the same length connecting $\mu_{j,n}^{+}$ to $\mu_{j+1,n}^{-}$ through full markings of $S - \tau_{j,n}$. Since $\sigma_{j,n} \subset \text{base}(\mu_{j,n}^{+}), \text{base}(\mu_{j+1,n}^{-})$, from (4.7) we conclude that all the markings intersect γ_n nontrivially, and this gives a bound on $d_{\gamma_n}(\mu^+_{j,n}, \mu^-_{j+1,n})$, as was desired.

Now we continue to prove Claim 4.9, note that

$$\varphi_{i,n}(\mu^-_{i,n}) = \mathcal{T}_{i,n}(\mu_i)$$

Thus, after applying $\varphi_{i,n}$ to all curves in the expression (4.6) in Claim 4.9 we get

$$d_\gamma(\mathcal{T}_{i,n}(\mu_i), \mu_i)$$

Now μ_i is fixed and contains γ as well as a transversal for γ. By Theorem 4.5 (4) $\mathcal{T}_{i,n}$ contains an arbitrarily large power of $\mathcal{D}_{\gamma_n}$, thus the claim follows.

Claims 4.7, 4.8 and 4.9 combine together with the triangle inequality and imply that for any $\gamma_n = \varphi^{-1}_{i,n}(\gamma)$ such that $\gamma \in \sigma_i - \hat{\tau}$ for some $i > 0$

$$d_{\gamma_n}(\mu(\zeta_n(0)), \mu(\zeta_n(T))) \to \infty \tag{4.8}$$

$$\text{as } n \to \infty$$

Now we continue to prove our theorem.

Proof of (1):

First we claim that after possibly passing to a subsequence there is $1 \leq i \leq k$ and $\gamma \in \sigma_i$ such that $\alpha_n = \varphi^{-1}_{i,n}(\gamma)$.

We prove the claim by contradiction, passing to a subsequence we may assume that J_n converge to a subinterval J. Note that the length of all J_n are uniformly bounded below, to see this note that by 1(a) $\ell_{\alpha_n}(\zeta_n(t))$ achieves the value ϵ_0 in each J_n and by 1(b) its infimum over J_n goes to 0 as $n \to \infty$. So by the convergence of lengths J also has a positive length.

Now recall the partition $0 = t_0 < t_1 < ... < t_{k+1}$, simplicies σ_i and $\hat{\tau}$ from the limiting picture. We claim that $k > 0$ and in fact for some $1 \leq i \leq k$ $t_i \in J$. If not, then for some $i > 0$ J is contained within (t_i, t_{i+1}) and so $\hat{\zeta}(J) \subset \mathcal{S}(\hat{\tau})$. Now by 1(b) $\inf_{t \in J_n} \ell_{\alpha_n}(\zeta_n(t)) \to 0$ as $n \to \infty$, thus after possibly passing to a subsequence $\alpha_n = \varphi^{-1}_{i,n}(\beta)$ for some $\beta \in \hat{\tau}$, but this implies that α_n has length going to 0 at every point of $\zeta_n(J_n)$ which contradicts 1(a).

Now by the above claim (1) follows from (4.8).

Proof of (2):

First we claim that after possibly passing to a subsequence there is $0 \leq i \leq k+1$ and $\gamma \in \sigma_i$ such that $\alpha_n = \varphi^{-1}_{i,n}(\gamma)$.

We prove the claim by contradiction, fix n and let $\alpha_n \notin \tau_{i,n}$ then $\alpha_n \pitchfork \text{base}(\mu_{i,n})$ for every $i = 0, 1, ..., k+1$, thus α_n intersects $\mu^-_{i,n}$ for every $i = 1, ..., k+1$ and $\mu^+_{i,n}$ for every $i = 0, ..., k$, moreover assume that $\alpha_n \notin \cup_{i=0,...,k+1}\sigma_{i,n}$ then since for each i, $\mu^+_{i,n} = \mathcal{T}_{i,n}(\mu^-_{i,n})$ and $\mathcal{T}_{i,n} \in \text{tw}(\sigma_{i,n})$

$$d_{\alpha_n}(\mu^-_{i,n}, \mu^+_{i,n})$$

is bounded for every $i = 1, ..., k$, moreover this bound does not depend on the choice of α_n as above.

As we saw in Claim 4.8 for every $j = 0, ..., k+1$

$$d_{\alpha_n}(\mu^+_{j,n}, \mu^-_{j+1,n})$$

is bounded and again the bound does not depend on the choice of α_n as above. Combing the above bounds by the triangle inequality

$$d_{\alpha}(\mu^-_{0,n}, \mu^+_{k,n})$$

is bounded for all n where the bound does not depend on the choice of α.

Now given a sequence α_n after possible passing to a subsequence either it is equal to $\varphi^{-1}_{i,n}(\gamma)$ for some $\gamma \in \sigma_i$ the we are done, if not by the above bound we have a bound for relative twist which contradicts 2(b) and the our claim follows.

By the above claim we need to show that for any γ_n

$$\inf_{t\in[0,T]} \ell_{\gamma_n}(\zeta_n(t)) \to 0$$

Consider the surface $\zeta_n(t_i)$ and the curve $\gamma_n = \varphi^{-1}_{i,n}(\gamma)$ such that $\gamma \in \sigma_i$ for some $i = 0, 1, ..., k+1$. Applying $\varphi_{i,n}$ we have

$$\ell_{\gamma_n}(\zeta_n(t_i)) = \ell_{\gamma}(\varphi_{i,n}(\zeta_n(t_i)))$$

Moreover, by Theorem 4.5 (3) we have $\varphi_{i,n}(\zeta_n(t_i)) \to \hat{\zeta}(t_i)$. But $\ell_\gamma(\hat{\zeta}(t_i)) = 0$, thus continuity of length-functions implies that $\ell_{\gamma_n}(\zeta_n(t_i)) \to 0$ as $n \to \infty$.

This finishes the proof of (2). $\square$

Remark 4.10. Fix a pants decomposition P and let $\big(\ell_\gamma, \theta_\gamma\big)_{\gamma\in P}$ be the corresponding Fenchel-Nielsen coordinate, let $x \in \mathrm{Teich}(S)$ be a point with $\theta_\alpha = \theta_0$ for some $\alpha \in P$, let $\epsilon_n \to 0$, consider the following two sequences of WP geodesic segments:

1. Let x_n be point with all coordinates equal to that of x, except $\theta_\alpha = \theta_0 + 2\pi\sqrt{-1}n$ and $\ell_\alpha(x_n) = \epsilon_n$. Then the WP geodesic segment $[x, x_n]$ has a uniformly bounded length and $d_\alpha(\mu(x), \mu(x_n)) \to \infty$.
2. Let x_n be point with all coordinates equal to that of x, except $\theta_\alpha = \theta_0$ and $\ell_\alpha(x_n) = \epsilon_n$. Then the WP geodesic segment $[x, x_n]$ has a uniformly bounded length and $d_\alpha(\mu(x), \mu(x_n)) = 0$.

These two simple examples show that there is no a-priori relation between a sequence of α_n-length-function going to zero at the end points of a sequence of WP geodesic segments and the $A(\alpha_n)$ subsurface coefficient of Bers markings at the end points of the geodesic segments. Thus the above theorem is sharp.

Note that in Theorem 4.6 (1) for positive constants s and ϵ_0, by the conditions 1(a) and 1(b) and the mean value theorem applied to the α_n-length-functions for n sufficiently large there is $t^*_n \in J_n$ such that $\dot{\ell}_{\alpha_n}(\zeta_n(t^*_n)) >$

$\frac{\epsilon_0}{T_n - 2s} \geq \frac{\epsilon_0}{T-2s}$. Now from the convexity of length-functions (3.1) $\ddot{\ell}_{\alpha_n}(\zeta_n(t)) > 0$ thus by the maximum principle

$$\ell_{\alpha_n}(\zeta_n(t)) > \frac{\epsilon_0}{T-2s}(t - t_n^*)$$

over the interval $[0, T_n]$.

Therefore, since $t_n^*, T - t_n^* \geq s$, we get a definite lower bound for the α_n-length-functions at both end points of ζ_n ($\zeta_n(T_n)$ and $\zeta_n(0)$).

We continue to prove two corollaries which are somewhat quantified versions of Theorem 4.6. These corollaries provide us with a kind of length-function versus twist parameter bounds over uniformly bounded length WP geodesic segments which will be used extensively in § 6.

Corollary 4.11. *(large twist $\Longrightarrow$ short curve) Given positive constants T, ϵ_0 and $\epsilon < \epsilon_0$ there is N with the following property:*

Let $\zeta : [0, T'] \to \mathrm{Teich}(S)$ be a WP geodesic segment of length $T' \leq T$ such that

1. $\sup_{t \in [0,T']} \ell_\gamma(\zeta(t)) \geq \epsilon_0$, *and*
2. $d_\gamma(\mu(\zeta(0)), \mu(\zeta(T'))) > N$

Then

$$\inf_{t \in [0,T']} \ell_\gamma(\zeta(t)) \leq \epsilon$$

Proof. The proof is by contradiction. Assume that the corollary does not hold. Then there is a sequence of WP geodesic segments $\zeta_n : [0, T_n] \to \mathrm{Teich}(S)$ parametrized by arc-length with length $T_n \leq T$ and $\gamma_n \in \mathcal{C}(S)$, such that

1. $\sup_{t \in [0,T_n]} \ell_{\gamma_n}(\zeta_n(t)) \geq \epsilon_0$ for every n,
2. $\inf_{t \in [0,T_n]} \ell_{\gamma_n}(\zeta_n(t)) \geq \epsilon$ for every n and
3. $d_{\gamma_n}(\mu(\zeta_n(0)), \mu(\zeta_n(T_n))) \to \infty$ as $n \to \infty$.

hold together. But this contradicts Theorem 4.6 (2).

□

Corollary 4.12. *(short curve $\Longrightarrow$ large twist)*

Given positive constants ϵ_0, T, N and s with $T > 2s$, there is ϵ with the following property:

Let $\zeta : [0, T'] \to \mathrm{Teich}(S)$ be a WP geodesic segment parametrized by arc-length of length $2s \leq T' \leq T$, $\gamma \in \mathcal{C}_0(S)$ and $J \subset [s, T' - s]$ be a subinterval such that:

1. $\sup_{t \in J} \ell_\gamma(\zeta(t)) \geq \epsilon_0$, *and*
2. $\inf_{t \in J} \ell_\gamma(\zeta(t)) \leq \epsilon$

Then

$$d_\gamma(\mu(\zeta(0)), \mu(\zeta(T'))) > N$$

Proof. The proof is again by contradiction. Assume that the corollary does not hold. Then there is a sequence of WP geodesic segments $\zeta_n : [0, T_n] \to \text{Teich}(S)$ parametrized by arc-length of length $2s \leq T_n \leq T$, $\gamma_n \in \mathcal{C}(S)$, subintervals $J_n \subset [s, T_n - s]$ and $\epsilon_n \to 0$, such that

1. $\sup_{t \in J_n} \ell_{\gamma_n}(\zeta_n(t)) \geq \epsilon_0$ for every n,
2. $\inf_{t \in J_n} \ell_{\gamma_n}(\zeta_n(t)) \leq \epsilon_n \to 0$ as $n \to \infty$ and
3. $d_{\gamma_n}(\mu(\zeta_n(0)), \mu(\zeta_n(T_n))) \leq N$ for every n.

hold together. But this contradicts Theorem 4.6 (1). □

5 Stable hierarchy paths

In this section we characterize all hierarchy paths which are *stable* in the pants graph.

Definition 5.1. ($D-$Stable subset) Given a function $D : \mathbb{R}^{\geq 1} \times \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ a subset $\mathcal{Y}$ of a metric space $\mathcal{X}$ is called $D-$stable if for any $K \geq 1$ and $C \geq 0$ a $(K, C)-$quasi-geodesic h with end points on $\mathcal{Y}$ be contained in the $D(K, C)$ neighborhood of $\mathcal{Y}$. Moreover, we call D the quantifier function of stability.

Brcok and Msure in [BM08] show that for $\xi(S) = 3$ $P(S)$ is strongly relatively hyperbolic with respect to the quasi-flats corresponding to separating curves.

Behrstock, Drutu and Mosher in [BDM09] study thick metric spaces, these are metric spaces with rank at least 2 where any two quasi-flats are connected through a chain of quasi-flats where any two consecutive quasi-flats in the chain has coarse intersection of infinite diameter. They show that thick spaces fail to be relatively hyperbolic with respect to any collection of quasi-flats. They observe that $P(S)$ in $\xi(S) > 3$ is a thick metric.

Let ρ be a hierarchy path with ending data (ν^-, ν^+), restricting subsurfaces with large subsurface coefficient $d_Y(\nu^-, \nu^+)$ to non-separated subsurfaces turns out to be equivalent to stability. Heuristically these hierarchy paths avoid quasi-flats in the pants graph corresponding to separating multi-curves on the surface. This is a generalization of stability of the hierarchy paths with bounded combinatorics ending data in the pants graph was proved in [BMM11].

To be able to save consider amount of work using results in the context of $\Sigma-$hulls and present our results in a more general setting we prove the stability for $\Sigma-$hulls. In §5.2 we prove that for any $\epsilon > 0$ sufficiently large a $\Sigma_\epsilon-$hull with A-narrow ending data is $D-$stable. Then the stability of hierarchy paths with narrow ending data will follow from the fact that they have a uniformly bounded Hausdorff distance from any $\Sigma_\epsilon-$hull with the same ending data. The Hausdorff distance depends only on A and ϵ. We prove the fact about the uniform bound on the Hausdorff distance in §5.1.

5.1 Narrow ending data

In this subsection we introduce the notion of *A-narrow ending data*. Then we will show that any hierarchy path with narrow ending data and any $\Sigma_\epsilon-$hull with the same ending data have uniformly bounded Hausdorff distance in the pants graph of the surface, where the Hausdorff distance depends on A and ϵ.

Definition 5.2. (Non-separated subsurface) A connected subsurface $Z \subseteq S$ is called non-separated if any connected component of $S - Z$ be either an annulus or a three holed sphere.

If Z is not connected or $S - Z$ has connected components other than annuli and three holed spheres then we call Z a *separated* subsurface.

Definition 5.3. ($A-$narrow) A pair of partial markings or laminations (μ^-, μ^+) is called $A-$narrow if every non-annular subsurface $Z \subseteq S$ with the property that

$$d_Z(\mu^-, \mu^+) > A$$

be a non-separated, non-annular subsurface of S.

A $\Sigma-$hull with $A-$narrow ending data is called an $A-$narrow $\Sigma-$hull. Similarly, a hierarchy with $A-$narrow ending data is called an $A-$narrow hierarchy.

In the rest of this subsection we show that a $\Sigma-$hull and any hierarchy path with the same narrow ending data are within a uniformly bounded Hausdorff distance of each other in the pants graph. We fix the constant $M = \max\{M_1, M_2, B_0\}$.

Theorem 5.4. *($\Sigma-$hull of narrow ending data)*

Given $\epsilon > M$ and $A > 2M + 2\epsilon$ there is $\Delta = \Delta(A, \epsilon)$ such that the Hasudorff distance of a $\Sigma_\epsilon - hull$ and a hierarchy path ρ with the same $A-$narrow ending data (ν^-, ν^+) is less than Δ.

Proof. Let $\rho : [m, n] \to P(S)$ and $i \in [m, n]$ then for every non-annular domain $Y \subseteq S$ Theorem 2.11 (6) says that there is $x_Y \in \text{hull}_Y(\nu^-, \nu^+)$ such that $d_Y(\rho(i), x_Y) \le M_2 \le \epsilon$. Thus from the definition of $\Sigma_\epsilon-$hull and since $\epsilon > M_2$ we have $\rho(i) \in \Sigma_\epsilon(\nu^-, \nu^+)$, consequently

$$|\rho| \subset \Sigma_\epsilon(\nu^-, \nu^+)$$

We continue by the following lemma:

Lemma 5.5. *There is $d = d(A, \epsilon)$ with the property that given $P \in \Sigma_\epsilon(\nu^-, \nu^+)$ there is $j \in [m, n]$ such that for every non-annular $Y \subseteq S$ we have*

$$d_Y(\rho(j), P) \le d \tag{5.1}$$

Proof. Let $e = A + M + 2\bar{\delta} + \epsilon + 1$. Define $E_Y = E_Y(P)$ to be the closure of the set $\{i \in [m, n] : d_Y(P, \rho(i)) \le e\}$, then E_Y is a connected subinterval of $[m, n]$.

If we show that $\bigcap_{Y \subseteq S} E_Y \neq \emptyset$, then for any $j \in \bigcap_{Y \subseteq S} E_Y$, $\rho(j)$ would satisfy (5.1) for every non-annular $Y \subseteq S$ and we are done.

Our strategy is to show that for any two subsurfaces Y and W

$$E_Y \cap E_W \neq \emptyset \tag{5.2}$$

then Helly's theorem in one dimension (see [Eck93]) implies that the intersection of all the intervals $\bigcap_{Y \subseteq S} E_Y$ is nonempty. We continue to prove (5.2) by considering several cases.

In the case that

$$d_Y(\nu^-, \nu^+) \leq A$$

then by hyperbolicity of $\mathcal{C}(Y)$ $\mathrm{diam}_Y(\mathrm{hull}_Y(\nu^-, \nu^+)) \leq A + 2\bar{\delta} + 1$ where $\bar{\delta} = \max_{Y \subseteq S} \delta_Y$. Moreover, $P \in \Sigma_\epsilon(\nu^-, \nu^+)$ thus we have

$$d_Y(P, \mathrm{hull}_Y(\nu^-, \nu^+)) \leq \epsilon$$

Moreover, by Theorem 2.11 (6) for every $i \in [m, n]$

$$d_Y(\rho(i), \mathrm{hull}_Y(\nu^-, \nu^+)) \leq M_2$$

The above three bounds and the triangle inequality imply that

$$d_Y(\rho(i), P) \leq e$$

Thus in this case $E_Y = [m, n]$ and obviously intersects any other E_W. Therefore, in the rest of proof we assume that

$$d_Y(\nu^-, \nu^+) > A$$

Consider the following collection of subsurfaces:

- $\mathcal{D} = \mathcal{D}(\nu^-, \nu^+) = \{$ non-annular $Y \subseteq S : d_Y(\nu^-, \nu^+) > A\}$

Then given $Y, W \in \mathcal{D}$ the $A-$narrow condition implies that either $Y \pitchfork W$, $Y \supsetneq W$ or $Y \subsetneq W$.

Note that by Theorem 2.11 (4) (monotonicity) and since $A > 2M$ for each $Y \in \mathcal{D}$ there are $i_Y^-, i_Y^+ \in J_Y$ such that

- $i_Y^- = \max\{i \in [n, m] : d_Y(\rho(i), \nu^-) \leq M\}$ and
- $i_Y^+ = \min\{i \in [n, m] : d_Y(\rho(i), \nu^+) \leq M\}$.

moreover $i_Y^- \leq i_Y^+$. Note that by Theorem 2.11 (4) and (5) if $i \leq i_Y^-$ then $d_Y(\rho(i), \nu^-) \leq M$, also if $i \geq i_Y^+$ then $d_Y(\rho(i), \nu^+) \leq M$.

We continue to discuss the above cases.

Case 1: $Y \pitchfork W$;

Without loss of generality assume that $Y < W$ (the case $W < Y$ can be treated exact same way). Then from Proposition 2.14 we have the following two pairs of inequalities:

(i) $d_Y(\nu^-, \partial W) > M$ and $d_W(\nu^-, \partial Y) \leq M$

(ii) $d_W(\nu^+, \partial Y) > M$ and $d_Y(\nu^+, \partial W) \leq M$

Note that in each pair of inequalities by the Behrstock's inequality, see Theorem 2.6, the first inequality implies the second one.

Here we claim that if $Y < W$ then we can choose i_Y^+ and i_W^- such that $i_Y^+ \leq i_W^-$. To see this note that by Theorem 2.11 (5) $d_Y(\rho(i), \nu^+) \leq M_2$ for every $i \in J_W$, now since $i_W^- \in J_W$ then $d_Y(\rho(i_W^-), \nu^+) \leq M_2$. Thus we pick a vertex on g_Y such that $v \leq i_W^-$, now let $i \in J_Y$ be such that by Theorem 2.11 (4) $\pi_Y(\rho(i_Y^+)) < v$ as vertices on g_Y.

We will discuss the following three cases and in each verify that

$$E_W \cap E_Y \neq \emptyset$$

Case 1.1: $d_Y(P, \nu^-) \leq M + \epsilon$;

We claim that

Claim 5.6. $E_Y \supseteq [m, i_Y^-]$

To see this note that if $i \leq i_Y^-$ then $d_Y(\rho(i), \nu^-) \leq M$, thus by the triangle inequality we have

$$d_Y(P, \rho(i)) \leq 2M + \epsilon$$

This implies that $i \in E_Y$, because $2M + \epsilon \leq A + M + \epsilon < e$ (recall that by assumption $A > 2M + 2\epsilon$) and the claim follows.

Now we claim that:

Claim 5.7. $E_W \supseteq [m, i_W^-]$.

To see this note that since $Y \in \mathcal{D}$ then $d_Y(\nu^-, \nu^+) > 2M + 2\epsilon$, moreover from inequality (ii) $d_Y(\partial W, \nu^+) \leq M$, also in this case $d_Y(P, \nu^-) \leq M + \epsilon$. Combining the three bounds by the triangle inequality we get $d_Y(P, \partial W) > \epsilon > M$, so by Behrstock's inequality we have

$$d_W(\partial Y, P) \leq M \tag{5.3}$$

Now from inequality (i) $d_W(\partial Y, \nu^-) \leq M$, which combined with (5.3) by the triangle inequality implies that

$$d_W(P, \nu^-) \leq 2M \leq M + \epsilon$$

Now as we saw in Claim 5.6 the last inequality implies that $E_W \supseteq [m, i_W^-]$ and the claim follows.

Moreover, we have $i_Y^- \leq i_W^-$ which together with the above two claims implies that $E_W \cap E_Y \neq \emptyset$.

Case 1.2: $\min\{d_Y(P, \nu^-), d_Y(P, \nu^+)\} > M + \epsilon$;

We claim that $E_Y \cap [i_Y^-, i_Y^+] \neq \emptyset$. To see this note that since $P \in \Sigma_\epsilon(\nu^-, \nu^+)$ there $y \in \operatorname{hull}_Y(\nu^-, \nu^+)$ such that $d_Y(y, P) \leq \epsilon$. Moreover, by Theorem 2.11 (6) there is $i \in [m, n]$ such that $d_Y(\rho(i), y) \leq M_2$. Thus by the triangle inequality $d_Y(P, \rho(i)) \leq M + \epsilon$ and since $M + \epsilon < A + M + \epsilon < e$ then $i \in E_Y$.

Moreover, by the triangle inequality $d_Y(\rho(i),\nu^-) > M$, therefore $i > i_Y^-$.

Similarly, we can show that $i < i_Y^+$. Thus $i \in [i_Y^-, i_Y^+]$ and the claim follows.

Now we claim that $E_W \supseteq [m, i_W^-]$. To see this note that from inequality (ii) $d_Y(\partial W, \nu^+) \leq M$, moreover in this case $d_Y(P,\nu^+) > M+\epsilon$ thus by the triangle inequality we get $d_Y(P,\partial W) > \epsilon > M$. Therefore, by Behrstock's inequality $d_W(P,\partial Y) \leq M$, which as we saw in Claim 5.7 implies our claim.

Now since $i_Y^+ < i_W^-$ by the above assertions about E_Y and E_W we have $E_Y \cap E_W \neq \emptyset$.

Case 1.3: $d_Y(P,\nu^+) \leq M+\epsilon$;

An argument similar to Case 1.1 implies that $E_Y \supseteq [i_Y^+, n]$, moreover since $Y < W$ Theorem 2.11 (5) implies that $J_W \subseteq E_Y$.

Moreover, it is immediate from definition that $J_W \cap E_W \neq \emptyset$. Thus $E_W \cap E_Y \neq \emptyset$

Case 2: $W \subsetneq Y$;

Recall the constant $e = A + M + 2\bar{\delta} + \epsilon + 1$.

Case 2.1: $d_Y(\partial W, P) \leq e - 1$;

By Theorem 2.11 (1) for every $i \in J_W$ $\partial W \subset \rho(i)$, thus for every $i \in J_W$ we have

$$d_Y(\rho(i), P) \leq e$$

which implies that $J_W \subset E_Y$.

Moreover, $J_W \cap E_W \neq \emptyset$. Thus $E_W \cap E_Y \neq \emptyset$.

Case 2.2: $d_Y(\partial W, P) > e - 1$;

By Theorem 2.11 (6) the Hausdorff distance in $\mathcal{C}(Y)$ of g_Y and $\mathrm{hull}_Y(\nu^-,\nu^+)$ is less than M_2, moreover since $P \in \Sigma_\epsilon(\nu^-,\nu^+)$ then $d_Y(P, \mathrm{hull}_Y(\nu^-,\nu^+)) \leq \epsilon$. Let x'_Y be a nearest points on g_Y to $\pi_Y(P)$. Then by the triangle inequality $d_Y(x'_Y, P) \leq M + \epsilon$.

Now let h be a geodesic connecting $\pi_Y(P)$ to x'_Y, we claim that h does not intersect the $1-$neighborhood of ∂W. Otherwise, ∂W would have distance at most

$$d_Y(P, x'_Y) + 1 \leq 1 + (M+\epsilon)$$

to $\pi_Y(P)$, see Figure 1 . But this contradicts the lower bound in assumption of this case.

By the above claim ∂W intersects every vertex of h so Bounded geodesic image theorem, Theorem 2.4, implies that

$$d_W(P, x'_Y) \leq G \tag{5.4}$$

In the rest of discussion let v be the initial vertex of g_Y in $\mathcal{C}(Y)$ and w be its final vertex.

Case 2.2.1: $v < x'_Y < \partial W$ as vertices on g_Y.



Figure 1: **Case 2.2:** If h intersects the $1-$neighborhood of ∂W then the distance in $\mathcal{C}(Y)$ between $\pi_Y(\partial W)$ and $\pi_Y(P)$ would be less than the assumption of this case.

Since g_Y is a tight geodesic then ∂W intersects every vertex of g_Y between v and x'_Y then by Theorem 2.4 $d_W(x'_Y, v) \leq G$ combining this inequality, (5.4) and Theorem 2.11 (5) by the triangle inequality we get

$$d_W(P, \nu^-) \leq 2M$$

Note that M_2 is a constant larger than $\xi(S)G$.

Now we claim that $E_W \supseteq [m, i^-_W]$. To see this note that if $i \in [m, i^-_W]$ then $d_W(\rho(i), \nu^-) \leq M$, thus by the triangle inequality $d_W(P, \rho(i)) \leq 3M + \epsilon$ now since $A > 2M$ the claim follows.

Moreover, by Theorem 2.11 (4) (monotonicity) there is $i \in E_Y$ such that $i < i^-_W$. Thus $E_W \cap E_Y \neq \emptyset$.

Case 2.2.2: $x'_Y < \partial W < w$ as vertices on g_Y.

In this case similar to the previous case we can show that $d_W(P, \nu^+) \leq M$.

Which again as the previous case implies that $E_W \supseteq [i^+_W, n]$. Moreover, there is $i \in E_W$ such that $i > i^+_W$. Thus $E_W \cap E_Y \neq \emptyset$.

In summary in all the above cases we verified that $E_Y \cap E_W \neq \emptyset$, thus as we explained from Helly's theorem in dimension one the lemma follows. □

Now we continue to prove the bound on the Hausdorff distance.

By the lemma inequality (5.1) holds for every non-annular $Y \subseteq S$ then the distance formula (2.3) implies that

$$d(P, \rho(j)) \leq C(d)$$

where $C(d)$ is the additive term in the distance formula corresponding to the threshold d.

This bound and the inclusion of $|\rho|$ in $\Sigma_\epsilon(\nu^-, \nu^+)$ together imply that the Hausdorff distance of ρ and $\Sigma_\epsilon(\nu^-, \nu^+)$ is bounded by $\Delta = \max\{\epsilon, C(d)\}$. Note that since d depends on A and ϵ, Δ also would depend on A and ϵ.

□

5.2 Stability in the pants graph

The following theorem is the main result of this subsection

Theorem 5.8. *(Stable hierarchy resolution path) Given $A > 0$ there is a quantifier function D such that any $A-$narrow hierarchy path is $D-$stable in the pants graph. Moreover, given D there is A' such that if a hierarchy path is $D-$stable then it is A'-narrow.*

First note that by Theorem 5.4 given narrow ending data (ν^-, ν^+) the $\Sigma_\epsilon-$hull ($\epsilon > F$) and a hierarchy path with the same ending data are within a bounded Hausdorff distance. Thus the stability of hull implies that of the hierarchy path. Thus we need to show that the $\Sigma-$hull is stable.

Our strategy to prove stability of the $\Sigma-$hull is to show that the projection $\Pi : P(S) \to \Sigma_\epsilon(\nu^-, \nu^+)$ ($\epsilon > F$) defined in Theorem 2.15 has the following contraction property:

Definition 5.9. (Contraction property)

Given positive constants R, B and $\eta \leq 1$ a subset $\mathcal{Y}$ of a metric space $\mathcal{X}$ is $(R, B, \eta)-$contracting if there is a map $\Pi : \mathcal{X} \to \mathcal{Y}$ with the following property:

For every $x, y \in \mathcal{X}$ if $d(x, \Pi x) > R$ then

$$d(x, y) \leq \eta d(x, \Pi y) \Longrightarrow d(\Pi x, \Pi y) \leq B$$

Then the standard Morse Lemma argument, see [MM99] Lemma 7.1 and [BH99] § III.H.1, implies that there is a quantifier function D depending only on R, B and η such that $\mathcal{Y}$ with contraction property is $D-$stable in $\mathcal{X}$.

The following proposition is a consequence of an elementary property of projections on geodesics in $\delta-$hyperbolic spaces which will be used often in the proof of the contraction property of $\Sigma-$hulls, see Lemma 5.12.

Proposition 5.10. *(Tree-like)*

Given $Y \subseteq S$, there are δ'_Y and K_Y with the following properties:

Let $\Pi : P(S) \to \Sigma_\epsilon(\nu^-, \nu^+)$ be the projection on $\Sigma-$hull for some $\epsilon > F$. Then for any $P, Q \in P(S)$ if $d_Y(\Pi P, \Pi Q) > K_Y$ then

$$d_Y(P, \Pi P) + d_Y(\Pi P, \Pi Q) + d_Y(\Pi Q, Q) \leq d_Y(P, Q) + \delta'_Y \tag{5.5}$$

Proof. The proposition follows from the following tree like property in $\delta-$hyperbolic spaces, see [Mah10], Propositions (3.1)-(3.4) for the proof.

Given points p, q and a geodesic σ in a $\delta-$hyperbolic space, let p' be a closet point projections of p on σ and q' be a closest point projection of q on σ. Then if $d(p', q') \geq 16\delta$ we have

$$d(p, p') + d(p', q') + d(q', q) \leq d(p, q) + 24\delta$$

Now by Theorem 2.15 in any subsurface $Y \subseteq S$ we have:

$$d_Y(x_Y, \Pi P) \leq F$$

where x_Y is a nearest point to $\pi_Y(P)$ on $\mathrm{hull}_Y(\nu^-, \nu^+)$.

Therefore, the triangle inequality implies that lemma holds for $K_Y = 16\delta_Y + 2F$ and $\delta'_Y = 24\delta_Y + 4F$. ☐

Recall that δ_Y the hyperbolicity constant of $\mathcal{C}(Y)$ ($Y \subseteq S$) depends only on the topological type of Y, also the constants δ'_Y and K_Y in the previous proposition. Moreover, subsurfaces of S have only finitely many different topological types, thus we can consider the following maximums over subsurfaces:

- $\bar{\delta} = \max_{Y \subseteq S} \delta_Y$, $\bar{K} = \max_{Y \subseteq S} K_Y$, and $\delta' = \max_{Y \subseteq S} \delta'_Y$.

Theorem 5.11. *(Narrow hulls are contracting)*
There is a positive constant $\bar{A}$ only depending on the topological type of S with the following property: Given $A \geq \bar{A}$ and $\epsilon > F$ there are positive constants R, B and $\eta \leq 1$ such that any $\Sigma_\epsilon-$hull with $A-$narrow ending data has the contraction property.

Proof. We prove that the projection $\Pi : P(S) \to \Sigma_\epsilon(\nu^-, \nu^+)$ defined in Theorem 2.15 has the contraction property.

Rather than proving the contraction property of the projection to $\Sigma-$hulls with narrow ending data we prove the following contrapositive of the contraction property as is stated in Definition 5.9:

- There are positive constants R, B and $\eta \leq 1$ such that
  If $d(P, \Pi P) > R$ then

$$d(\Pi P, \Pi Q) > B \Longrightarrow d(P, Q) > \eta d(P, \Pi P)$$

Lemma 5.12. *There is $\bar{A}$ only depending on the topological type of S with the property that for any $A \geq \bar{A}$ there is $\delta(A)$ such that a $\Sigma_\epsilon-$hull ($\epsilon > F$) with $A-$narrow ending data has the property that for every $Z \subseteq S$*

$$d_Z(P, Q) > d_Z(P, \Pi P) - \delta \tag{5.6}$$

Let us first see how this lemma implies the contrapositive of the contraction property.

Assume that (5.6) holds for every $Z \subseteq S$ then by the distance formula (2.3) we have

$$\begin{aligned} d(P,Q) &\asymp_{K,C} \sum_{\substack{Z\subseteq S\\ \text{non-annular}}} \{d_Z(P,Q)\}_A \\ &> \sum_{\substack{Z\subseteq S\\ \text{non-annular}}} \{d_Z(P,\Pi P)-\delta\}_A \end{aligned}$$

Recalling the definition of $\{.\}_A$, for any term in the last sum above we have

$$\{d_Z(P,\Pi P)-\delta\}_A \geq \frac{A}{A+\delta}\{d_Z(P,\Pi P)\}_{A+\delta}$$

Moreover, for the threshold $A_1 = A+\delta$ the distance formula (2.3) gives us

$$\sum_{\substack{Z\subseteq S\\ \text{non-annular}}} \{d_Z(P,\Pi P)\}_{A+\delta} \asymp_{K_1,C_1} d(P,\Pi P)$$

Therefore, we get

$$d(P,Q) > \eta' d(P,\Pi P) - c$$

where $\eta' = \frac{A}{K(A+\delta)K_1}$ and $c = \frac{AC_1}{K(A+\delta)} + C$.

Now let R be large enough such that $\eta = \eta' - \frac{c}{R} > 0$. Then for any $P \in P(S)$ with the property that $d(P,\Pi P) > R$ we have

$$\begin{aligned} d(P,Q) &> \eta'(d(P,\Pi P)) - c = [(\eta' - \frac{c}{R})d(P,\Pi P)] - c + (\frac{c}{R})d(P,\Pi P) \\ &> \eta d(P,\Pi P) \end{aligned}$$

This shows that for the constants R and η as above the projection $\Pi : P(S) \to \Sigma_\epsilon(\nu^-,\nu^+)$ has the contrapositive contraction property.

Now we continue to prove the stated lemma.

Proof of Lemma 5.12. First we choose $\bar{A}$ as follows:

$$\bar{A} = \bar{K} + G + M_2 + 2F \tag{5.7}$$

Here $\bar{K} = \max_{Y\subseteq S} K_Y$, G is the bound from the Bounded geodesic image theorem, Theorem 2.4, M_2 is the constant from Theorem 2.11 (5). Moreover, constant F comes from Theorem 2.15.

We fix $A > \bar{A}$ and let (ν^-,ν^+) be an $A-$narrow ending data. Also fix $A' = A + 2\Delta(A,\epsilon) + 2M_2$ as the threshold in the distance formula. Here $\Delta(A,\epsilon)$ is the constant from Theorem 5.4.

To begin note that if

$$d_Z(P,\Pi P) \leq A' + 2(F_1+4)$$

Then for $\delta = A' + 2(F_1 + 1)$, (5.6) holds. Thus in the rest of proof we assume that Z is a subsurface with

$$d_Z(P, \Pi P) > A' + 2(F_1 + 1)$$

We need to discuss the following two cases:

Caes 1: $d_Z(\Pi P, \Pi Q) > A'$;
Then since $A' > \bar{K} > K_Z$ (5.5) implies that

$$d_Z(P, Q) > d_Z(P, \Pi P) - \delta'$$

where $\delta' = \max_{W \subseteq S} \delta'_W$. Thus (5.6) holds for $\delta = \delta'$.

Case 2: $d_Z(\Pi P, \Pi Q) \leq A'$;
In this case Z has no contribution to $d(\Pi P, \Pi Q)$. But still if we let $B = K' + C'$, K' and C' are the constants in the distance formula corresponding to the thresholds A', then

$$d(\Pi P, \Pi Q) > K' + C'$$

thus by the distance formula (2.3) there is a subsurface $W \subseteq S$ such that

$$d_W(\Pi P, \Pi Q) > A'$$

Therefore in the rest of proof we can assume that the following three conditions on subsurface coefficients of subsurfaces Z and W hold

(i) $d_Z(P, \Pi P) > A' + 4(F_1 + 4)$,

(ii) $d_Z(\Pi P, \Pi Q) \leq A'$ and

(iii) $d_W(\Pi P, \Pi Q) > A'$

and continue to show that there is δ depending on A such that inequality (5.6) holds for Z.

Note that by Condition (iii)

$$d_W(\Pi P, \Pi Q) > A'$$

Now by Theorem 5.4 there are $\widehat{\Pi P}$ and $\widehat{\Pi Q}$ on a hierarchy path ρ with the ending data (ν^-, ν^+) such that for $\Delta = \Delta(A, \epsilon)$,

$$d(\Pi P, \widehat{\Pi P}) \leq \Delta \text{ and } d(\Pi P, \widehat{\Pi P}) \leq \Delta$$

thus by (2.2)

$$d_W(\Pi P, \widehat{\Pi P}) \leq \Delta \text{ and } d_W(\Pi P, \widehat{\Pi P}) \leq \Delta$$

Therefore by the triangle inequality we have

$$d_W(\Pi P,\Pi Q) > A' - 2\Delta$$

Now by Theorem 2.11 (5) we have

$$d_W(\nu^-,\nu^+) > A' - 2\Delta - 2M_2 > A$$

Now since the ending data (ν^-,ν^+) is $A-$narrow the subsurface $W \subseteq S$ is a non-separated subsurface, which excludes the case that W and Z are disjoint.

Thus we need to discuss the following three cases: $W \pitchfork Z$, $W \subsetneq Z$ and $Z \subsetneq W$. Note that $W = Z$ is essentially Case 1 which was already discussed.

Case 2.1: $W \pitchfork Z$;
In this case, let

- $k = k(P) = \lfloor \frac{d_Z(P,\Pi P)}{F_1+4} \rfloor$ and
- $k' = \lfloor \frac{A'}{F_1+4} \rfloor$.

By the choice of k

$$d_Z(\Pi P, P) > k(F_1+4) \tag{5.8}$$

Moreover, by Condition (i) we have

$$k > k' + 2$$

We claim that

$$d_Z(\Pi P,\partial W) \leq (k'+2)(F_1+4) \tag{5.9}$$

Otherwise,

$$d_Z(\Pi P,\partial W) > (k'+2)(F_1+4)$$

Then using ΠP to define a partial order, that is let $x_Y = \pi_Y(\Pi P)$ in Definition 2.9 we have

$$Z \prec_{k'+2} W \tag{5.10}$$

Moreover, by Condition (iii) and since $A' > 2(F_1+4)$

$$W \ll_2 \Pi Q \tag{5.11}$$

Now (5.10), (5.11) and the transitivity property of $\ll$, Theorem 2.10 (2), imply that

$$Z \ll_{k'+1} \Pi Q$$

therefore,

$$d_Z(\Pi Q, \Pi P) > (k'+1)(F_1+4) > A'$$

But this contradicts Condition (ii) and our claim follows. Therefore, in the rest of this case we can assume that (5.9) holds.

Now (5.8), (5.9) and the triangle inequality in $\mathcal{C}(Z)$ imply that

$$d_Z(P, \partial W) > (k-k'-2)(F_1+4) \tag{5.12}$$

Now using P to define a partial order, that is let $x_Y = \pi_Y P$ in Definition 2.9, (5.12) implies that

$$Z \prec_{k-k'-2} W \tag{5.13}$$

Moreover, by Condition (iii)

$$d_W(\Pi P, \Pi Q) > A'$$

Now since $A' > \bar{K} > K_W$ (5.5) implies that for $\delta' = \max_{Y \subseteq S} \delta'_W$

$$\begin{aligned} d_W(P,Q) &> d_W(\Pi P, \Pi Q) - \delta' \\ &> A' - \delta' \end{aligned}$$

Thus we have

$$W \ll_m Q \tag{5.14}$$

for $m = \lfloor A' - \delta' \rfloor \geq 2$.

(5.13), (5.14) and the transitivity property of $\ll$ imply that

$$Z \ll_{k-k'-3} Q$$

Therefore,

$$\begin{aligned} d_Z(P,Q) &> (\lfloor \frac{d_Z(P,\Pi P)}{(F_1+4)} \rfloor - k' - 3)(F_1+4) \\ &> d_Z(P,\Pi P) - (k'+4)(F_1+4) \end{aligned}$$

which implies that inequality (5.6) holds for $\delta = (k'+4)(F_1+4)$ in Case 2.1.

Case 2.2: $W \subsetneq Z$;

Let $H(P, \Pi P)$ be a hierarchy between P and ΠP. By Condition (i) and since $A' > M_1$ Z is a component domain of $H(P, \Pi P)$.

Let x be a nearest point to $\pi_Z(Q)$ on $\mathrm{hull}_Z(P, \Pi P)$ and let h be a geodesic in $\mathcal{C}(Z)$ connecting $\pi_Z(Q)$ to x, see Figure 2 the left diagram. Moreover, let T

be a slice of $H(P,\Pi P)$ such that $d_Z(T,x) \leq M_2$, note that Theorem 2.11 (6) guarantees that such a T exists.

Now since T is a slice of $H(P,\Pi P)$ by Theorem 2.11 (6) there is $y \in \mathrm{hull}_W(P,\Pi P)$ such that

$$d_W(T,y) \leq M_2 \tag{5.15}$$

Now by the definition of projection Π and Condition (iii) we have

$$d_W(Q,\mathrm{hull}_W(P,\Pi P)) > d_W(\Pi P,\Pi Q) - 2F$$

Thus for $y \in \mathrm{hull}_W(P,\Pi P)$ as above we have

$$d_W(Q,y) > d_W(\Pi P,\Pi Q) - 2F \tag{5.16}$$

Now (5.15) and (5.16) combined by the triangle inequality imply that

$$\begin{aligned} d_W(Q,T) &> d_W(Q,y) - d_W(y,T) \\ &> d_W(\Pi P,\Pi Q) - 2F - M_2 \\ &> A' - 2F - M_2 > G \end{aligned} \tag{5.17}$$

where the last inequality holds by Condition (iii) and since $A' > \bar{A}$, see the choice of $\bar{A}$ in (5.7).

We claim that ∂W is in the $1-$neighborhood of h in $\mathcal{C}(Z)$ that is

$$d_Z(\partial W,h) \leq 1 \tag{5.18}$$

Otherwise, ∂W intersects every vertex of the geodesic h in $\mathcal{C}(Z)$ which connects $\pi_Z(T)$ to $\pi_Z(Q)$ and by the Bounded geodesic image theorem, Theorem 2.4, $\pi_W(h)$ would have length less than G in $\mathcal{C}(W)$ which contradicts (5.17) above and the claim follows, see Figure 2 the right diagram.

Now we claim that $\pi_Z(\partial W)$ is in the $1-$neighborhood of any geodesic $k \subset \mathrm{hull}_Z(\nu^-,\nu^+)$ connecting $\pi_Z(\nu^-)$ to $\pi_Z(\nu^+)$. Otherwise, ∂W would intersect every vertex of k and Theorem 2.4 would imply that $d_W(\nu^-,\nu^+) \leq G$. This contradicts Condition (iii) and the claim follows. Thus by the claim we have

$$d_Z(\partial W,k) \leq 1 \tag{5.19}$$

(5.18), (5.19) and the triangle inequality in $\mathcal{C}(Z)$ imply that h and k have distance less than 2 in $\mathcal{C}(Z)$, thus

$$d_Z(x,\Pi P) \leq 2 + F \tag{5.20}$$

Considering the triangle $\triangle(PxQ)$ in $\mathcal{C}(Z)$ the thin triangle property and (5.20) imply that



Figure 2: **Case 1.2**, ∂W is in the $1-$neighborhood of geodesics h and $\text{hull}_Z(\nu^-, \nu^+)$, thus $d_Z(x, \Pi P) \leq 2 + F$

$$d_Z(P, Q) > d_Z(P, \Pi P) - (2) - \bar{\delta} - F$$

where $\bar{\delta} = \max_{Y \subseteq S} \delta_Y$.

This shows that the inequality (5.6) holds for $\delta = 2 + \bar{\delta} + F$.

Case 2.3: $Z \subsetneq W$;

We claim that ∂Z is in the $1-$neighborhood of any geodesic $k \subset \text{hull}_W(P, \Pi P)$ connecting $\pi_W(P)$ to $\pi_W(\Pi P)$, that is

$$d_W(\partial Z, k) \leq 1 \tag{5.21}$$

To see this note that in this case $\partial Z \in \mathcal{C}(W)$ thus if (5.21) does not hold ∂Z would intersect every vertex of k then Theorem 2.4 implies that $d_Z(P, \Pi P) \leq G < A'$ but this contradicts Condition (i).

Now let $l \subset \text{hull}_W(Q, \Pi Q)$ be any geodesic connecting $\pi_W(Q)$ to $\pi_W(\Pi Q)$, see Figure 3 the left diagram. Condition (iii), (5.21) and the definition of Π imply that

$$d_W(\partial Z, l) > A' - 2F - 1 > 1$$

therefore ∂Z intersects every vertex of l, so again Theorem 2.4 implies that

$$d_Z(Q, \Pi Q) \leq G$$

Now considering the quadrilateral $\square(PQ\Pi P\Pi Q)$ in $\mathcal{C}(Z)$ since $d_Z(Q, \Pi Q) \leq G$, see Figure 3 the right digram, the thin triangle property implies that

$$d_Z(P, Q) > d_Z(P, \Pi P) - F - \bar{\delta}$$



$\mathcal{C}(W)$ $\mathcal{C}(Z)$

Figure 3: **Case 2.3**, ∂Z is in $1-$neighborhood of $\text{hull}_W(\Pi P, \Pi Q)$, moreover $\pi_W(\Pi P)$ and $\pi_W(\Pi Q)$ have distance greater than $A' \geq 2$ in $\mathcal{C}(W)$. Thus ∂Z intersects every vertex of a geodesic connecting Q to ΠQ, therefor the bounded geodesic image theorem implies that $d_Z(Q, \Pi Q) \leq G$.

which shows that the inequality (5.6) holds for $\delta = F + \bar{\delta}$.

Therefore the lemma follows by choosing δ to be the maximum of δ's worked out in the above cases. This finishes the proof of Lemma 5.12. $\square$

Finishing the proof of Theorem 5.11: Recall the discussion before Lemma 5.12 for R, η depending on A and as we saw in the lemma $B = K' + C'$ the $\Sigma_\epsilon-$hull ($\epsilon > F$) with $A-$narrow ending data ($A > \bar{A}$) has the contraction property. $\square$

Remark 5.13. Note that as $A \to \infty$

$$R \to \infty,\ \eta \to 0 \text{ and } B \to \infty$$

thus applying the Morse lemma argument we get $\Sigma-$hulls and hierarchy paths with worse and worse stability property, that is the corresponding D goes to ∞, see Definition 5.1.

Proof Theorem 5.8. We fix $\epsilon = \max\{M, F\}$ then Theorem 5.8 for every $A > \bar{A}$ gives us a uniform contraction for every $A-$narrow $\Sigma-$hull. Moreover Theorem 5.4 gives us a uniform Hausdorff distance between a hierarchy path and a $\Sigma-$hull with the same $A-$narrow ending data. These two together imply that an $A-$narrow hierarchy path is stable with the quantifier function D depending only on A.

We continue to show that for given a function$D$ there is $A' > 0$ such that a $D-$stable hierarchy path ρ is $A'-$narrow. We prove its contrapositive, that is there is A' such that if ρ is not $A'-$narrow then it is not $D-$stable.

Fix a logarithmic spiral $\{x(t) = ae^{bt}\cos t,\ y(t) = ae^{bt}\sin t : t \in \mathbb{R}\}$ in the Euclidean plane, there are K_0 and C_0 for which this curve is a quasi-geodesic.

Given $D : \mathbb{R}^{\geq 1} \times \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ there is d such that for any path in the plane we can move the logarithmic spiral to put in a position such that intersects the path at two points with distance d and does not stay in the $D(K_0, C_0)$ neighborhood the path.

Moreover, by Theorem 2.11 (5) for $A' > d$ if an ending data (ν^-, ν^+) is not $A'-$narrow then there is a quasi-flat corresponding to the separated domain with subsurface coefficient larger than A' in which any hierarchy path ρ with ending data (ν^-, ν^+) and the logarithmic spiral can be put in the configuration described above. This shows that ρ is not $D-$stable.

□

5.3 Fellow-traveling

Definition 5.14. (Fellow-traveling) Given $D \geq 0$, we say that two parametrized paths $h : I \to \mathcal{X}$ and $h' : I' \to \mathcal{X}$, $D-$fellow-travel each as parametrized paths or simply $D-$fellow-travel each other if

1. For every $i \in I$ there is $i' \in I'$ such that $d(h(i), h'(i')) \leq D$ and vice versa. In other words the Hausdorff distance of $h(I)$ and $h'(I')$ are bounded by D.

2. The map assigning to each $i \in I$ any $h'(i')$ such that $d(h(i), h'(i')) \leq D$ be a reparametrization of h' as a quasi-geodesic with constants depending only on D and vice versa.

   Moreover, given parametrized path h and h' and interval $J \subset I$ we say that h $D-$fellow-travel h' over J if there is $J' \subset I'$ such that $h|_J$ $D-$fellow-travel $h'|_{J'}$ as above.

Theorem 5.15. *Given a positive constant A there is $D = D(A)$ with the property that any WP geodesic segment $g : [a, b] \to \mathrm{Teich}(S)$ with $A-$narrow ending data $D-$fellow travels a hierarchy path ρ with the same ending data.*

Proof. By Theorem 5.8 ρ is $D-$stable for a function D depending on A. Moreover by Theorem 3.3 (Brock's quasi-isometry) $Q(g)$ is a $(K_{\mathrm{WP}}, C_{\mathrm{WP}})-$quasi geodesic in $P(S)$. Thus any $Q(g)$ with end points on $|\rho|$ stays in the $D(K_{WP}, C_{WP})$ neighborhood of $|\rho|$ in $P(S)$. Moreover, ρ is a $(K, C)-$quasi-geodesic in $P(S)$ with constants depending only on the topological type of S.

Now $Q(g)$ stays in the $D(A)$ neighborhood of ρ and both $Q(g)$ and ρ are quasi-geodesic with constants depending on the topological type of S, therefore ρ and $Q(g)$ fellow-travel each other as parametrized paths in the pants graph. □

Let $Q(g)$ and ρ $D-$fellow-travel each other as parametrized quasi-geodesics in the pants graph. For given $i \in [m, n]$ take the smallest interval containing every $t \in [a, b]$ such that $d(Q(g(t)), \rho(i)) \leq D$. Then define $N_{\rho,g}(i) \subset [a, b]$ to be the $K_{\mathrm{WP}}(2D + 1) + C_{\mathrm{WP}}$ neighborhood of the this interval.

$\bigcup_{i\in[m,n]} N_{\rho,g}(i)$ covers $[a,b]$, thus $N_{\rho,g}$ defines a correspondence from parameters in $[m,n]$ to uniformly bounded length subintervals of $[a,b]$, moreover we have

Proposition 5.16. *There are constants $K \geq 1$ and $C \geq 0$ depending only on D and the the topological type of S such that for any $i,j \in [m,n]$*

$$d_{Haus}(N(i),N(j)) \asymp_{K,C} |i-j|$$

where d_{Haus} is the Hausdorff distance of the intervals $N(i)$ and $N(j)$ as subsets of $\mathbb{R}$.

By Theorem 5.15 this proposition in particular applies to a WP geodesic segment with $A-$narrow ending data and a hierarchy path with the same ending data. We say that a Weil-Petersson geodesic $g : [a,b] \to \overline{\mathrm{Teich}(S)}$ parametrized by arc-length $D-$fellow-travels a hierarchy resolution path $\rho : I \to P(S)$ if $Q(g)$ $D-$fellow-travels ρ in the pants graph.

6 Itinerary of a Weil-Petersson geodesic segment

Itinerary of a WP geodesic g in Teichmüller space refers to list of short curves (curves with length bounded by a given sufficiently small ϵ), the time along g over which each curve is short and the order in which the short curves appear along g.

In this section we present our results on the control of length-functions and twist parameters along WP geodesics with narrow ending data.

Our main result asserts that on a suitably shrunk subinterval of an interval over which all subsurface coefficients except some annular subsurfaces whose core curves consist a non-separating multi-curve (boundary of a non-separated subsurface) are bounded the length of the core curves are arbitrary small.

Recall the parameter correspondence map N for the parameters of hierarchy path and a fellow-traveling WP geodesic from §5.3.

Theorem 6.1. *(short curve)*

Given positive constants A, R, R' and an ϵ sufficiently small, there is a constant $\bar{s} = \bar{s}(A,R,R',\epsilon)$ with the following property:

Let $g : [a,b] \to \mathrm{Teich}(S)$ be a WP geodesic segment with $A-$narrow ending data.

Let $\rho : [m,n] \to P(S)$ be a hierarchy with the same ending data, assume that a non-separated domain Z has $(R,R')-$bounded combinatorics over $[i_1,i_2] \subset J_Z$, moreover let $t_1 \in N(i_1)$ and $t_2 \in N(i_2)$.

Then if $t_2 - t_1 \geq 2\bar{s}$ we have

$$\ell_{\partial Z}(g(t)) \leq \epsilon$$

for every $t \in [t_1+\bar{s}, t_2-\bar{s}]$.

Moreover, in Lemma 6.8 (2) we establish a lower bound for the length of all the curves which are not in the multi-curve (boundary of Z), this lower bound holds on a much longer subinterval containing the subinterval where the multi curve is short. Theorem 1.5 in the introduction follows from these two bounds.

6.1 Isolated annular subsurfaces

In this subsection we prove some extra combinatorial lemmas which together with the fellow-traveling theorem for hierarchy paths with narrow ending data, Theorem 5.15, give us a combinatorial frame work in which we will be able to control length-functions along WP geodesics.

Definition 6.2. (Non-annular bounded combinatorics over an interval) Given $R > 0$, let $\rho : [m, n] \to P(S)$ be a hierarchy path, then we say that a component domain Z has non-annular $R-$bounded combinatorics over an interval $[i_1, i_2] \subset J_Z \subset [m, n]$ if for any essential, proper, non-annular subsurface $Y \subsetneq Z$ we have

$$d_Y(\rho(i_1), \rho(i_2)) \leq R$$

Definition 6.3. (Bounded combinatorics over an interval) Given $R, R' > 0$, let $\rho : [m, n] \to P(S)$ be a hierarchy path, then we say that a component domain Z has $(R, R')-$bounded combinatorics over an interval $[i_1, i_2] \subset J_Z \subset [m, n]$ if for any essential, proper, non-annular subsurface $Y \subsetneq Z$ we have

$$d_Y(\rho(i_1), \rho(i_2)) \leq R$$

moreover, for any essential, annular subsurface with core curve $\gamma \in \mathcal{C}(Z)$ we have

$$d_\gamma(\rho(i_1), \rho(i_2)) \leq R'$$

Remark 6.4. Recall the notion of $R-$bounded combinatorics hierarchy path with ending data (ν^-, ν^+) in [BMM11] where for every proper subsurface $Y \subsetneq S$

$$d_Y(\nu^-, \nu^+) \leq R$$

The condition in Definition 6.3 is a local version of this condition, because we bound subsurface coefficients between two pants decomposition along the hierarchy path rather than the ending data.

Moreover, we bound the subsurface coefficients of the proper subsurfaces of a subsurface Z which is not necessarily S, moreover we have independent bounds R' and R respectively for annular subsurfaces and non-annular subsurfaces.

In the following lemma we show that over a subintrval of a hierarchy path where a non-separated subsurface Z has non-annular bounded combinatorics the subsurface projection π_Z gives an effective reparametrization of g_Z, see Theorem 2.11 (3), more precisely in this case $\pi_Z \circ \rho$ is a reparametrization of g_Z as a quasi-geodesic.

Lemma 6.5. *Given a positive constant R there are constants $K_R \geq 1$ and $C_R \geq 0$ with the following property:*

Let $\rho : [m,n] \to P(S)$ be a hierarchy path and let Z be a non-separated, non-annular domain with non-annular $R-$bounded combinatorics over $[i_1, i_2] \subset [m,n]$. Then for any $i, j \in [i_1, i_2]$

$$d(\rho(i), \rho(j)) \asymp_{K_R, C_R} d_Z(\rho(i), \rho(j))$$

Proof. By the distance formula (2.3),

$$d(\rho(i), \rho(j)) \asymp_{K,C} \sum_{\substack{Y \subseteq S \\ \text{non-annular}}} \{d_Y(\rho(i), \rho(j))\}_A$$

Now since $i, j \in J_Z$ by Theorem 2.11 (1) $\partial Z \subset \rho(i)$ and $\partial Z \subset \rho(j)$. Therefore, for any $Y \pitchfork Z$

$$d_Y(\rho(i), \rho(j)) \leq 2$$

Moreover, since Z has complement consisting of only annuli and three holed spheres there is no Y disjoint from Z contributing to the above sum.

Also since $i, j \in [i_1, i_2] \subset J_Z$, Theorem 2.11 (4) and the non-annular $R-$bounded combinatorics of Z over $[i_1, i_2]$ for any $Y \subsetneq Z$ imply that

$$d_Y(\rho(i), \rho(j)) \leq d_Y(\rho(i_1), \rho(i_2)) + 2M_2 \leq R + 2M_2$$

The above bounds on the subsurface coefficients contributing to the sum imply that if we raise A to $A_R > \max\{2M_2 + R, 2\}$ we get

$$d(\rho(i), \rho(j)) \asymp_{K_R, C_R} d_Z(\rho(i), \rho(j))$$

as was desired, where K_R and C_R are the constants corresponding to the threshold A_R in the distance formula. $\square$

Let ρ and g $D-$fellow-travel each other and $N = N_{g,\rho}$ be the parameter correspondence from §5.3. Given $i, j \in [m,n]$ let $r \in N(i)$ and $s \in N(j)$, then by $D-$fellow-traveling $d(\rho(i), Q(g(r))) \leq D$ and $d(\rho(j), Q(g(s))) \leq D$, thus by (2.2) for every non-annular subsurface $Y \subseteq S$

$$d_Y(\rho(i), Q(g(r)) \leq D \text{ and } d_Y(\rho(j), Q(g(s))) \leq D$$

so

$$d_Y(\rho(i), \rho(j)) \asymp_{1,2D} d_Y(Q(g(r)), Q(g(s)))$$

But such a comparison does not exit for annular subsurfaces, so we consider *isolated annular subsurfaces* along hierarchy paths. In Lemma 6.7 we prove a comparison for the subsurface coefficient of an isolated annular subsurface along ρ and a $D-$fellow-traveling WP geodesic which is uniform in D.

Definition 6.6. (Isolated annular subsurface) Let $\rho : [m.n] \to P(S)$ be a hierarchy path. We say that an annular subsurface $A(\gamma)$ with core curve γ is $(w, r)-$isolated at $i \in [m, n]$, if there is a pants decomposition $\widehat{Q}$ such that $\gamma \in \widehat{Q}$ and $d(\widehat{Q}, \rho(i)) \leq r$, moreover there are non-separated, non-annular subsurfaces Z_1 and Z_2 with $\gamma \notin \partial Z_1$ and $\gamma \notin \partial Z_2$ such that Z_1 and Z_2 have non-annular $R-$bounded combinatorics over intervals $[i_1, i_1 + w] \subset J_{Z_1}$ and $[i_2 - w, i_2] \subset J_{Z_2}$ respectively, where $i_1 + w < i_2 - w$, for the illustration of isolation see Figure 4.

Lemma 6.7. *(Annular coefficient comparison) Given positive constants D, r and R there are constants $w = w(D, r, R)$ and $B = B(D)$ with the following properties:*

Let $\rho : [m, n] \to P(S)$ be a hierarchy path, let $g : [a, b] \to \mathrm{Teich}(S)$ be a WP geodesic parametrized by arc-length $D-$fellow-traveling ρ and let $t_1 \in N(i_1)$ and $t_2 \in N(i_2)$.

If $A(\gamma)$ be $(w, r)-$isolated at i, then for any two partial markings μ_1, μ_2 with $\mathrm{base}(\mu_1) = Q(g(t_1))$ and $\mathrm{base}(\mu_1) = Q(g(i_2))$, we have

$$d_\gamma(\mu_1, \mu_2) \asymp_{1,B} d_\gamma(\rho(i_1), \rho(i_2)) \tag{6.1}$$

Moreover,

$$\min\{\ell_\gamma(g(t_1)), \ell_\gamma(g(t_2))\} \geq \omega(L_S) \tag{6.2}$$

Proof. There is $\widehat{Q}$ with $\gamma \in \widehat{Q}$ and $d(\widehat{Q}, \rho(i)) \leq r$, moreover by the $D-$fellow-traveling $d(\rho(i_1), Q(g(t_1))) \leq D$, thus (2.2) implies that for every non-annular subsurface $Y \subseteq S$ we have that $d_Y(\widehat{Q}, \rho(i)) \leq r$ and $d_Y(Q(g(t_1)), \rho(i+w)) \leq D$. In particular, for Z_1, the domain with non-annular $R-$bounded combinatorics on left, we have the following bounds:

(*) $d_Z(\widehat{Q}, \rho(i)) \leq r$ and $d_Z(P, \rho(j)) \leq D$.

Let $w(D, R) = K_R(D + r + C_R + 2)$, where K_R and C_R are the constants in Lemma 6.5.

Now let P' be a pants decomposition on a geodesic in $P(S)$ connecting $\rho(i_1)$ to $Q(g(t_1))$ then

$$\begin{aligned} d_{Z_1}(P', \widehat{Q}) &> d_{Z_1}(\rho(i_1), \rho(i)) - d_{Z_1}(\widehat{Q}, \rho(i)) - d_{Z_1}(\rho(i_1), P') \\ &> \frac{1}{K_R} d(\rho(i), \rho(i+w)) - C_R - r - D \\ &> 2 \end{aligned}$$

The second inequality follows from the bounds in (*) and Lemma 6.5 and the triangle inequality, see Figure 5. The last inequality follows from the choice of w.

Now since $\gamma \in \widehat{Q}$ and $\gamma \notin \partial Z_1$ the above inequality implies that $\gamma \pitchfork P'$. Thus, the diameter of the projection to $A(\gamma)$ of the geodesic in $P(S)$ connecting $\rho(i_1)$ to $Q(g(t_1))$ is bounded above by D, that is $d_\gamma(Q(g(t_1)), \rho(i_1)) \leq D$. Now μ_1 is a partial markings with $\mathrm{base}(\mu_1) = Q(g(t_1))$ and $\gamma \pitchfork \mathrm{base}(\mu_1)$ then from the definition of projection of markings to annular subsurfaces $d_\gamma(\rho(i_1), \mu_1) \leq$



Figure 4: Isolated annular subsurface: non-annular, non-separated domains Z_1 and Z_2 have non-annular bounded combinatorics respectively on the left and the right blue subintervals of length at least w, moreover $\gamma \in \widehat{Q}$, so $A(\gamma)$ is $(w,r)-$isolated at i.

$D+2$. The same argument on the right hand side and for the domain Z_2 gives us $d_\gamma(\rho(i_1),\mu_1) \leq D+2$. Now by the triangle inequality (6.1) follows for $B = 2(D+2)$.

As we saw above $\gamma \pitchfork Q(g(t_1))$ and $\gamma \pitchfork Q(g(t_2))$, now since each curve in both $Q(g(t_1))$ and $Q(g(t_2))$ has length at most the Bers constant L_S the lower bound (6.2) for the length of γ follows form the Collar lemma.

□

6.2 Length-function control

Corollaries 4.11 and 4.12 in §4 give us a control on the development of the Dehn twist about a curve versus the change of its length-function along WP geodesic segments. The control is uniform in the length of the geodesic segment and the supremum of the length-function along the geodesic segment.

(6.1) in Lemma 6.7 gives a comparison for the subsurface coefficient of an isolated annular subsurface along a hierarchy path and the subsurface coefficient of the partial markings at the end points of a WP geodesic segment fellow-traveling the hierarchy path. The comparison is uniform in the fellow-traveling distance D.

We use these technical tools in this subsection to prove our main results about the control of length-functions and twist parameters along WP geodesic segments with narrow ending data.

Lemma 6.8. *(Rough upper bound)*

Given positive constants D, R and R', there are constants $s = s(D,R)$, $\bar{\epsilon} = \bar{\epsilon}(D,R')$ and $l = l(D,R')$ with the following properties:

Let $\rho : [m,n] \to P(S)$ be a hierarchy path, assume that a non-separated domain Z has $(R,R')-$bounded combinatorics over $[i_1,i_2] \subset J_Z$.

Let $g : [a,b] \to \overline{\text{Teich}(S)}$ be a WP geodesic parametrized by arc-length which $D-$fellow travels ρ, moreover let $t_1 \in N(i_1)$ and $t_2 \in N(i_2)$.

Then if $t_2 - t_1 > 2s$

1. *$\ell_\alpha(g(t)) \leq l$ for every $\alpha \in \partial Z$ and*

2. $\ell_\gamma(g(t)) > \bar{\epsilon}$ *for every* $\gamma \notin \partial Z$

for every $t \in [t_1 + s, t_2 - s]$.

Proof. Given $r > 0$ assume that an annular subsurface $A(\gamma)$ is $(w, r)-$isolated at $i_0 \in [i_1, i_2]$ where $w = w(D, r, R)$ is the constant from Lemma 6.7 and Z is the non-annular bounded combinatorics domain on both sides. Let K and C be the constants for the parameter correspondence map $N_{\rho,g}$ form Proposition 5.16 and let $s = Kw + C$. Let $t_0 \in N(i_0)$, moreover pick $t_0^+ \in N(i_0 + w)$ and $t_0^- \in N(i_0 - w)$ such that $t_0^+ - t_0^- = 2s$.

By Lemma 6.7 we have

$$\min\{\ell_\gamma(g(t_0^-)), \ell_\gamma(g(t_0^-))\} \geq \omega(L_S) \tag{6.3}$$

and, for $B = B(D)$

$$d_\gamma(\mu(i_0 + w), \mu(i_0 - w)) \asymp_{1,B} d_\gamma(\mu(g(t_0^+)), \mu(g(t_0^-))) \tag{6.4}$$

Claim 6.9. There is $s' < s$ such that if $\inf_{t \in [t_0^-, t_0^+]} \ell_\gamma(g(t)) \leq \frac{\omega(L_S)}{2}$ then the infimum is realized in the interval $[t_0^- + s', t_0^+ - s']$.

We have $t_0^+ - t_0^- = 2s$ so the claim follows from the bound (6.3) and Corollary 3.6.

We start by establishing the lower bound (2) for the length of every $\gamma \notin \partial Z$, which is the following:

There are $s = s(D, R)$ and $\bar{\epsilon} = \bar{\epsilon}(D, R')$ such that for every $\gamma \notin \partial Z$

$$\ell_\gamma(g(t)) \geq \bar{\epsilon}$$

for every $t \in [t_1 + s, t_2 - s]$

Let $r = 0$ and $w = w(D, 0, R)$ and $s = Kw + C$ (K and C are the constants for the map N). Let $i_0 \in [i_1 + w, i_2 - w]$, $t_0 \in N(i_0)$, $t_0^- \in N(i_0 - w)$ and $t_0^+ \in N(i_0 + w)$ be as above.

We continue to show that there is a lower bound for the length of γ at $g(t_0)$.

If $\ell_\gamma(g(t_0)) \geq L_S$ then we already have the lower bound. Otherwise, $\gamma \in Q(g(t_0))$, thus $A(\gamma)$ is $(w, 0)$-isolated at i_0 where the non-annular bounded combinatorics domain on both sides is Z.

By Claim (6.9) and Corollary 4.12 there is $\bar{\epsilon} \leq L_S$ such that if

$$\inf_{t \in [t_0^-, t_0^+]} \ell_\gamma(g(t)) \leq \bar{\epsilon}$$

then

$$d_\gamma(\mu(g(t_0^-)), \mu(g(t_0^+))) > R' + 2M_2 + B \tag{6.5}$$

Thus (6.4) and (6.5) give us

$$d_\gamma(\mu(i_0 + w), \mu(i_0 - w)) > R' + 2M_2$$

But by Theorem 2.11 (6) this contradicts the bound on annular coefficients along $[i_1, i_2]$.

Therefore, we get the lower bound $\bar{\epsilon}$ for length of γ at $g(t_0)$. Now note that $t_0 \in N(i_0)$, also $i_0 \in [i_1 + w, i_2 - w]$ was arbitrary. Moreover, as we saw in §5.3 $\bigcup_{i \in [i_1+w, i_2-w]} N(i)$ covers $[t_1 + s, t_2 - s]$. Thus for $s = Kw + C$ we have $\ell_\gamma(g(t)) > \bar{\epsilon}$ for any $t \in [t_1 + s, t_2 - s]$ and the claim follows. Note that since $t_0^+ - t_0^- = 2s$ for any $t_0 \in [t_1 + s, t_2 - s]$ then $\bar{\epsilon}$ is uniform over this interval and does not depend on the value of the parameter t_0.

We continue to establish the rough upper bound (1). We choose a hyperbolic surface $x \in V_{L_S}(\rho(i_0))$ as follows:

(*) $\ell_\alpha(x) = L_S$ for every $\alpha \in \partial Z$ and $\frac{L_S}{2} \leq \ell_\alpha(x) \leq L_S$ for every $\alpha \in \rho(i_0) - \partial Z$, then $Q(x) = \rho(i_0)$,

(**) Let $d_0 = \min\{\mathbf{r}(L_S, \frac{L_S}{4}), \mathbf{r}(\frac{3L_S}{4}, \frac{L_S}{4})\}$ then by Corollary 3.6 there is x such that the bounds in (*) for the length of curves in $\rho(i_0)$ hold, and $d_{\mathrm{WP}}(g(t_0), x) \geq d_0$. To see this let $\alpha \in \rho(i_0) - \partial Z$ then if $\ell_\alpha(g(t_0)) \geq \frac{3L_S}{4}$ pick x with $\ell_\alpha(x) = \frac{L_S}{2}$, otherwise $\ell_\alpha(g(t_0)) \leq \frac{3L_S}{4}$ and we pick x with $\ell_\alpha(x) = L_S$.

note that d_0 does not depend on i_0,

(***) A Bers marking of x, which we denote it by μ_x, is equal to a marking slice of any hierarchy $H(\nu^-, \nu^+)$ which has base equal to $\rho(i_0)$, note that by changing the twist parameters of curves in $\rho(i_0)$ we can get such an x.

Note that there can be several such surfaces. By (*) x has a definite positive injectivity radius $\mathrm{inj}(x)$ only depending on L_S.

Let $[x, g(t_0)]$ be the WP geodesic connecting x to $g(t_0)$. By the $D-$fellow-traveling of ρ and g the length of u is bounded by $d_1 = K_{\mathrm{WP}} D + C_{\mathrm{WP}}$.

Let $u : [0, d] \to \mathrm{Teich}(S)$ be a parametrization of $[x, g(t_0)]$ by arc-length such that $u(0) = x$ and $u(d) = g(t_0)$ then $d_0 \leq d \leq d_1$.

To get a rough upper bound for lengths of boundary components of Z at $g(t_0)$ first we establish a lower bound for the length of every $\gamma \notin \partial Z$ along u, then the rough upper bound follows from a simple compactness argument.

Let $D' = K_{\mathrm{WP}} d_1 + C_{\mathrm{WP}}$ then D' bounds the distance between $Q(x) = Q(u(0))$ and any $Q(u(t))$ for $t \in [0, d]$.

In the rest of proof we let $r = D'$ and $w = w(D, D', R)$. As before fix $i_0 \in [i_1 + w, i_2 - w]$, assuming $[i_1, i_2]$ is long enough, and let $t_0 \in N(i_0)$, $t_0^- \in N(i_0 - w)$ and $t_0^+ \in N(i_0 + w)$, see Figure 5.

Here we make the following three choices:

(i) By Claim (6.9) and Corollary 4.12 there is $\epsilon_1 < L_S$ such that if

$$\inf_{t \in [t_0^-, t_0^+]} \ell_\gamma(g(t)) \leq \epsilon_1$$

then

$$d_\gamma(\mu(g(t_0^-)), \mu(g(t_0^+))) > R' + 2M_2 + B$$



Figure 5: The domain Z has $(R, R')-$bounded combinatorics over the red interval. y is the point along $[x, g(t)]$ where the infimum of the $\gamma-$length-function is realized thus $\gamma \in Q(y)$. These two together imply that γ is $(w, D')-$isolated at i_0.

(ii) Let $I \subseteq [t_0^-, t_0^+]$ be a subinterval with at least one end point t_0^- or t_0^+ then by Corollary 4.11 there is N_0 such that if

$$d_\gamma(\mu(g(t_0^-)), \mu(g(t_0^+))) > N_0$$

then

$$\inf_{t \in [t_0^-, t_0^+]} \ell_\gamma(g(t)) \leq \epsilon_1$$

(iii) Let $\bar{\epsilon} = \bar{\epsilon}(D, R)$ be the lower bound we established above for $\ell_\gamma(g(t_0))$ and every $\gamma \notin \partial Z$. Then by (**) and Corollary 3.6 there is s_0 with $2s_0 < d_0 < d$ such that if $r \in [d - s_0, d]$ then $\ell_\gamma(u(r)) > \frac{\bar{\epsilon}}{2}$ and if $r \in [0, s_0]$ then $\ell_\gamma(u(r)) > \mathrm{inj}(x)$.

Then let $\epsilon_2 < \min\{\mathrm{inj}(x), \frac{\bar{\epsilon}}{2}\}$ be such that from Corollary 4.12 if

(a) $\sup_{r \in [s_0, d - s_0]} \ell_\gamma(u(r)) \geq \min\{\mathrm{inj}(x), \frac{\bar{\epsilon}}{2}\}$, and

(b) $\inf_{r \in [s_0, d - s_0]} \ell_\gamma(u(r)) \leq \epsilon_2$

then

$$d_\gamma(\mu(u(0)), \mu(u(d))) > N_0 + R' + 2M_2 + D$$

We continue to rule out the possibility that $\inf_{r \in [0,d]} \ell_\gamma(u(t)) \leq \epsilon_2$ for some $\gamma \notin \partial Z$, where ϵ_2 is chosen in (iii).

The proof is by contradiction, assume that the infimum is less than ϵ_2, then by choice of s_0 in (iii) we may assume that the infimum is realized in $J = [s_0, d - s_0]$ that is

$$\inf_{r \in J} \ell_\gamma(u(r)) \leq \epsilon_2 \tag{6.6}$$

Now by the choice of ϵ_2 in (iii) and since $\mu_x = \mu(u(0))$ and $\mu(g(t_0)) = \mu(u(d))$

$$d_\gamma(\mu_x, \mu(g(t_0))) > N + R' + 2M_2 + D \tag{6.7}$$

Moreover, by (6.6) there is $y \in u(J)$ such that $\gamma \in Q(y)$.

Since the length of u is less than d_1 we have $d(Q(y), \rho(i_0)) \leq D'$ (recall that $D' = K_{\mathrm{WP}} d_1 + C_{\mathrm{WP}}$).

Let $\mu(i_0 - w)$ be any marking such that $\mathrm{base}(\mu(i_0 - w)) = \rho(i_0 - w)$, and let $\mu(g(t_0^-))$ be any Bers marking on $g(t_0^-)$ then as we saw in the proof of Lemma 6.7

$$d_\gamma(\mu(i_0 - w), \mu(g(t_0^-))) \leq D \tag{6.8}$$

Moreover, by (***) $\mathrm{base}(\mu_x) = \rho(i_0)$. Thus by the bound on annular coefficients over $[i_1, i_2]$ we have

$$d_\gamma(\mu_x, \mu(i_0 - w)) \leq R' + 2M_2 \tag{6.9}$$

Now (6.7), (6.8) and (6.9) combined by the triangle inequality imply that

$$d_\gamma(\mu(g(t_0)), \mu(g(t_0^-))) > N_0$$

Now the choice of N_0 in (ii) implies that $\inf_{t \in [t_0^-, t_0]} \ell_\gamma(g(t)) \leq \epsilon_1$. Thus by the choice of ϵ_1 in (i)

$$d_\gamma(\mu(g(t_0^-)), \mu(g(t_0^+))) > R' + 2M_2 + B \tag{6.10}$$

Moreover, $A(\gamma)$ is $(w, D')-$isolated at i_0 where the isolating domain on both sides is Z thus (6.4) holds, which together with (6.10) implies that

$$d_\gamma(\mu(i_0 + w), \mu(i_0 - w)) > R' + 2M_2$$

but this contradicts the bound on the annular coefficients over $[i_1, i_2]$. This contradiction shows that

$$\inf_{t \in [0,d]} \ell_\gamma(u(t)) > \epsilon_2$$

for every $\gamma \notin \partial Z$.

Therefore, we get the lower bound ϵ_2 for length of every $\gamma \notin \partial Z$ along u. Note that ϵ_2 does not depend on the value $i_0 \in [i_1 + w, i_2 - w]$ and $t_0 \in N(i_0)$, therefore is uniform along $[i_1 + w, i_2 - w]$.

Now consider the set of all WP geodesic segments u of length bounded by d_0 and d_1 emanating from x with the property that $\ell_\gamma(u(t)) \geq \epsilon_2$ for every $\gamma \notin \partial Z$ and every t in the domain of u. Assume that length of $\alpha \in \partial Z$ goes to ∞ in a sequence of segments u_k in this set, then since Teichmüller space is locally compact at x after possibly passing to a subsequence u_k converge to a WP geodesic segment u emanating from x of bounded by d_0 and d_1. Moreover, by continuity of length functions $\ell_\gamma(u(t)) \geq \epsilon_2$ for every $\gamma \notin \partial Z$ and every t in the domain of u. Now let $\hat{\gamma}$ be a curve in $\mathcal{C}(S)$ with minimum intersection number (1 or 2) with α, then by the Collar lemma $\ell_{\hat{\gamma}}(u_k(t)) \to 0$ as $k \to \infty$ which contradicts the lower bound we established above.

Thus, there is a bound l depending only on D and R' such that

$$\ell_{\partial Z}(g(t_0)) \leq l$$

Now $i_0 \in [i_1+w, i_2-w]$ and $t_0 \in N(i_0)$ were arbitrary, moreover $\bigcup_{i\in[i_1+w,i_2-w]} N(i)$ covers $[t_1+s, t_2-s]$, thus we get the desired rough upper bound (1) for every $t \in [t_1+s, t_2-s]$ where $s = Kw + C$.

□

Corollary 6.10. *(Recurrence) Given positive constants A, R and R' there are constants $s = s(A,R)$ and $\bar{\epsilon} = \bar{\epsilon}(A,R,R')$ with the following properties:*

Let $g : [a,b] \to \overline{\mathrm{Teich}(S)}$ be a WP geodesic segment parametrized by arc-length with $A-$narrow ending data.

If S has $(R,R')-$bounded combinatorics over $[t_1,t_2]$ where $t_2 - t_1 > 2s$, then

$$\mathrm{inj}(g(t)) \geq \bar{\epsilon}$$

for every $t \in [t_1+s, t_2-s]$.

Proof. Let $D = D(A)$ be the fellow-traveling distance of g and the hierarchy path with same ending data. Let S be the subsurface with $(R,R')-$bounded combinatorics then by Lemma 6.8 for every $\gamma \in \mathcal{C}(S)$ (the boundary of S consists of only peripheral curves) we have $\ell_\gamma(g(t)) \geq \bar{\epsilon}$ for every $t \in [t_1+s, t_2-s]$, which means $\mathrm{inj}(g(t)) \geq \bar{\epsilon}$ on this interval.

Note that for example the middle point of $g([t_1+s, t_2-s])$ projects to a recurrence point to the thick part of moduli space for the projection of g to the moduli space. □

Remark 6.11. Compare the recurrence corollary to the main result of [BMM11] which asserts that for every $R > 0$ there is $\epsilon > 0$ such that if the ending data of a WP geodesic g be $(R,R')-$bounded combinatorics then the geodesic is $\epsilon-$co-bounded in Teichmüller space, that is the geodesic projects to the $\epsilon-$thick part of moduli space.

In what follows we use an inductive argument which involves convexity properties of length-functions along WP geodesics and Wolpert's estimate for pairing of length-function gradients to sharpen the rough upper bound obtained in the previous lemma.

Proof of Theorem 6.1. We show that:

Given positive constants $l, \bar{\epsilon}$ and $\epsilon \leq \bar{\epsilon}$ if for a WP geodesic g the length-function bounds

1. $\ell_\alpha(g(t)) \leq l$ for every $\alpha \in \partial Z$, and
2. $\ell_\gamma(g(t)) \geq \bar{\epsilon}$ for every $\gamma \notin \partial Z$.

hold for every $t \in [t^-, t^+]$.

Then for any $\bar{s}$ sufficiently large if $t^+ - t^- > 2\bar{s}$ then for every $\alpha \in \partial Z$

$$\ell_\alpha(g(t)) \leq \epsilon$$

for every $t \in [t^- + \bar{s}, t^+ - \bar{s}]$.

The proof is by induction on $|\partial Z|$ the number of boundary components of Z.

Step of induction

Assume that for any non-separated subsurface Z' with $|\partial Z'| < |\partial Z|$ if the bounds (1) and (2) hold then the above conclusion holds.

The proof breaks into several steps.

Step 1: We show that given $\epsilon < \bar{\epsilon}$ for any $\bar{s}$ sufficiently large if $t^+ - t^- > 2\bar{s}$ then there is $\alpha \in \partial Z$ such that

$$\ell_\alpha(g(t)) \leq \epsilon$$

for some $t \in [t^- + \bar{s}, t^+ - \bar{s}]$.

First we claim that if $g|_{[t^-,t^+]}$ stays in the $\epsilon-$thick part of Teichmüller space then there is an upper bound T for length of the interval $[t^-, t^+]$.

Fix $\hat{\alpha} \in \partial Z$, by (1) above $\ell_{\hat{\alpha}}(g(t)) \leq l$ for every $t \in [t^-, t^+]$, thus by the mean-value theorem if $t^+ - t^- > \frac{l}{\epsilon}$ then there is $t^* \in [t^-, t^+]$ such that $|\dot{\ell}_{\hat{\alpha}}(g(t^*))| < \epsilon$. Moreover, since we have assumed that $g|_{[t^-,t^+]}$ is in the $\epsilon-$thick part $\ell_{\hat{\alpha}}(g(t^*)) > \epsilon$.

Now again since $g|_{[t^-,t^+]}$ is in the $\epsilon-$thick part then by Theorem 3.8 (3.1) there is $c(\epsilon) > 0$ for which the differential inequality $\ddot{\ell}_{\hat{\alpha}}(g(t)) \geq c(\epsilon)\ell_{\hat{\alpha}}(g(t))$ holds on the interval $[t^-, t^+]$, thus the maximum principle implies that

$$\ell_{\hat{\alpha}}(g(t)) \geq \ell_{\hat{\alpha}}(g(t^*)) \cosh\left(\sqrt{c(\epsilon)}(t - t^*))\right) + \dot{\ell}_{\hat{\alpha}}(g(t^*)) \sinh\left(\sqrt{c(\epsilon)}(t - t^*))\right)$$

Now by the choice of t^* the left hand side of the above inequality is a positive, convex function on $[t^-, t^+]$, moreover by the bound (2) above $\ell_{\hat{\alpha}}(g(t)) < l$ on this interval. So the right hand side is a positive, convex function bounded above by l over $[t^-, t^+]$, now it is immediate that there is an upper bound T depending only on l and ϵ for the length of the interval $[t^-, t^+]$.

The above claim implies that if $t^+ - t^- > T$ then there are curves which get shorter than ϵ at some time along $g|_{t^-,t^+]}$.

Moreover, by the bound (2) above component curves of ∂Z are the only curves which can get shorter than $\epsilon < \bar{\epsilon}$ along $g|_{[t^-,t^+]}$.

Thus, we conclude that if $t^+ - t^- > T$ then there is $t \in [t^-, t^+]$ and $\alpha \in \partial Z$ such that

$$\ell_\alpha(g(t)) \leq \epsilon$$

Step 2: We show that for any $L > 0$, if $t^+ - t^- > (2|\partial Z| + 1)L + 2T$ then there is $\alpha \in \partial Z$ such that $\ell_\alpha(g(t)) \leq \epsilon$ on a subinterval of $[t^-, t^+]$ of length at least L.

Consider a partition of $[t^- + T, t^+ - T]$ to $2|\partial Z| + 1$ subintervals each one of length at least L, then by Step 1 for each of these subintervals there is a time at which a component curve of ∂Z gets shorter that ϵ. Now by the pigeon hole



Figure 6: Step of induction: $\alpha \in \partial Z$ is shorter than ϵ' along $g|_{[t_\alpha^-, t_\alpha^+]}$, $g' : [t^-, t^+] \to \mathcal{S}(\alpha)$ connects x and y the projections of respectively $g(t_\alpha^-)$ and $g(t_\alpha^-)$ on $\mathcal{S}(\alpha)$. For $2l$ and $\frac{\bar{\epsilon}}{2}$ the length bounds (i) and (ii) hold for the subsurface $Z' = Z - \alpha$ along g'. Then by the assumption of induction the length of every $\alpha' \in \partial Z'$ gets shorter than $\frac{\epsilon}{2}$ along g' over a suitably shrunk subinterval of $[t^-, t^+]$.

principle there is $\alpha \in \partial Z$ such that $\ell_\alpha(g(t)) \leq \epsilon$ at some time in at least three of the partition subintervals. Therefore, by the convexity of length-functions

$$\ell_\alpha(g(t)) \leq \epsilon$$

on an interval of length at least L.

Step 3: We show that given $\epsilon < \bar{\epsilon}$ for any $\bar{s}$ sufficiently large if $t^+ - t^- > 2\bar{s}$ then $\ell_{\partial Z}(g(t)) \leq \epsilon$ at some $t \in [t^- + \bar{s}, t^+ - \bar{s}]$.

Fix $\epsilon' < \bar{\epsilon}$ which will be determined, let $L = \bar{s}' + 2\sqrt{2\pi\epsilon'}$, where $\bar{s}'$ is the following constant: Given $l, \frac{\epsilon}{2}$ and $2\bar{\epsilon}$ by the assumption of induction for the WP geodesic $g|_{[t^-, t^+]}$ we have that for any $\bar{s}'$ sufficiently large if $t^+ - t^- \geq 2\bar{s}'$ then

$$\ell_{\partial Z'}(g(t)) \leq \frac{\epsilon}{2}$$

for every $t \in [t^- + \bar{s}', t^+ - \bar{s}']$.

Now by Step 2 if $t_2 - t_1$ be sufficiently large there is $\alpha \in \partial Z$ such that $\ell_\alpha(g(t)) \leq \epsilon'$ on an interval of length at least L. Denote this interval by $[t_\alpha^-, t_\alpha^+]$.

Project $g(t_\alpha^-)$ and $g(t_\alpha^+)$ to the $\alpha-$stratum and denote the projection points respectively by x and y. Since the $\alpha-$stratum is geodesically convex and totally geodesic in $\overline{\text{Teich}(S)}$ there is a WP geodesic segment $g' : [t^-, t^+] \to \mathcal{S}(\alpha)$ parametrized by arc-length connecting x to y.

Since $\ell_\alpha(g(t_\alpha^-)) \leq \epsilon'$ and $\ell_\alpha(g(t_\alpha^+)) \leq \epsilon'$ then by Wolpert's estimate for the distance of a point to a stratum (Corollary 4.10 in [Wol08]) we have $d_{\text{WP}}(g(t_\alpha^-), x) \leq \sqrt{2\pi\epsilon'}$ and $d_{\text{WP}}(g(t_\alpha^+), y) \leq \sqrt{2\pi\epsilon'}$. Moreover, $\overline{\text{Teich}(S)}$ equipped with the WP metric is a CAT(0) space, therefore by the CAT(0) comparison (see [BH99]) the distance between any point on $g([t_\alpha^-, t_\alpha^+])$ and its nearest point on g' is less than $\sqrt{2\pi\epsilon'}$.

Now we choose ϵ' such that:

- $\sqrt{2\pi\epsilon'} \leq \min\{\mathbf{r}(\bar{\epsilon}, \frac{\bar{\epsilon}}{2}), \mathbf{r}(2l, l), \mathbf{r}(\epsilon, \frac{\epsilon}{2})\}$.

where **r** is the function from Corollary 3.6.

By the choice of ϵ' we have the following bounds for every $t \in [t^-, t^+]$

(i) $\ell_\alpha(g'(t)) \leq 2l$ for every $\alpha \in \partial Z$

(ii) $\ell_\gamma(g'(t)) \geq \frac{\bar{\epsilon}}{2}$ for every $\gamma \notin \partial Z$

Now since α is a non-separating curve the above two bounds and the assumption of induction implies that for every $\bar{s}'$ sufficiently large if $t^+ - t^- > 2\bar{s}'$ then

$$\ell_{\alpha'}(g'(t)) \leq \frac{\epsilon}{2}$$

for every $\alpha' \in \partial Z - \alpha$ and every $t \in [t^- + \bar{s}', t^+ - \bar{s}']$, see Figure 6.

Thus again by the choice of ϵ'

$$\ell_{\alpha'}(g(t)) \leq \epsilon$$

for every t in the interval along $g|_{[t_\alpha^-, t_\alpha^+]}$ consisting of the nearest points to $g'|_{[t^- + \bar{s}', t^- - \bar{s}']}$. By the CAT(0) comparison for $\bar{s}'' = \bar{s} + \bar{s}' + 2\sqrt{2\pi\epsilon'}$ we get the following bound for the length of every $\alpha' \in \partial Z - \alpha$

$$\ell_\alpha(g(t)) \leq \epsilon$$

for every $t \in [t_1 + \bar{s}'', t_2 - \bar{s}'']$.

Moreover, since $[t_1 + \bar{s}'', t_2 - \bar{s}''] \subset [t_\alpha^-, t_\alpha^+]$ the above bound holds on this interval for α as well.

This finishes the step of induction.

Base of induction

∂Z consists of only one non-separating curve α. Let $D = D(A)$ be the fellow-traveling distance from Theorem 5.15.

Let g $D-$fellow-travels ρ over $[i_1, i_2]$ where Z has $(R, R')-$bounded combinatorics, let $t_1 \in N(i_1)$ and $t_1 \in N(i_1)$. Then by Lemma 6.8 there are $l = l(D, R), \bar{\epsilon} = \bar{\epsilon}(D, R')$ and $s = s(D, R)$ with the property that for every $t \in [t_1 + s, t_2 - s]$

1. $\ell_\alpha(g(t)) \leq l$, and
2. $\ell_\gamma(g(t)) \geq \bar{\epsilon}$ for every $\gamma \neq \alpha$.

Now given any $\epsilon < \bar{\epsilon}(D, R')$ Step 1 of the step of induction and the convexity of the $\alpha-$length-function imply that for any $\bar{s} > 2T(l, \epsilon)$ if $t_2 - t_1 > 2\bar{s}$ then $\ell_\alpha(g(t)) \leq \epsilon$ for every $t \in [t_1 + \bar{s}, t_2 - \bar{s}]$.

□

7 Weil-Petersson Geodesics

In this section we use the control on length-functions along WP geodesic segments from §6 and the hierarchy paths with prescribed list of large domains from appendix to provide examples of Weil-Petersson geodesic rays with certain behavior in the moduli space, for example diverging rays and closed geodesics in the thin part of moduli space. We also provide a recurrence criterion for WP geodesics in terms of ending laminations.

In §7.1 by extracting limits of WP geodesic segments with ending data on a single infinite hierarchy path with narrow ending data we construct WP geodesics whose behavior mimic the combinatorial properties of hierarchy paths and have a prescribed itinerary.

In [PWW10], the authors construct examples of closed WP geodesics staying outside a given compact subset of moduli space in $\xi(S) = 1$. Jeff Brock has given examples of diverging WP geodesic rays with minimal filling ending lamination.

Both constructions start with a piecewise geodesic in the Weil-Petersson completion of Teichmüller space. Then applying high Dehn twists manage to replace it with a piecewise geodesic with arbitrary small exterior angles. Further a kind of Shadowing lemma gives a single diverging or a closed WP geodesic. In these examples the relation between the ending data and the itinerary of the ray is not explicit.

A motivation for these constructions is the results about the behavior of Teichmüller geodesics. Cheung and Masur in [CM06] give examples of diverging geodesic rays with uniquely ergodic vertical lamination. Examples of closed Teichmüller geodesics staying in the thin part of moduli space worked out by Hamenstädt in [Ham] and Rafi in [Raf].

7.1 Weil-Petersson geodesic rays with prescribed itinerary

Let ρ be an infinite hierarchy path with narrow ending data (ν^-, ν^+), then ν^+ is a minimal lamination filling a non-separated subsurface $Z \subseteq S$. Consider the WP geodesic segment connecting a base point x with $Q(x) = \nu^-$ in the interior of Teichmüller space to c_n the maximally nodal hyperbolic surface at $\rho(n)$. We denote this segment by $[x, c_n]$ and denote its parametrization by arc-length by r_n.

In the following lemma we show that after possibly passing to a subsequence r_n limit to an infinite geodesic ray in the WP visual sphere at x. We denote this infinite geodesic ray by r_ρ. As will be explained in Theorem 7.2 the behavior of r_ρ mimics the combinatorial properties of the hierarchy path ρ. In § 7.2, 7.3, 7.4 we use the rays with itinerary prescribed with ending data with a prescribed list of large domains to provide for examples of WP geodesics with a specific behavior in the moduli space.

Lemma 7.1. *(Infinite ray) Let $\rho : [0, \infty) \to P(S)$ be a hierarchy path with ending data (ν^-, ν^+), then after possibly passing to a subsequence the geodesic segments $[x, c_n]$ converge to an infinite ray in the Weil-Petersson visual sphere at x with forward ending lamination ν^+.*

Proof. The proof follows from almost the exact same line in [BM08] for the proof of the surjection of the weighted laminations (page 2482-84). Since ρ is infinite in the forward direction there is a subsurface Z such that $\nu^+ = \nu' \cup \partial Z$ where $\nu' \in \mathcal{EL}(Z)$ is determined by the only infinite geodesic of ρ in $\mathcal{C}(Z)$, moreover by the narrow condition Z is a non-separated subsurface.

By the non-refraction theorem, Theorem 3.2, each one of the WP geodesic segments $[x, c_n]$ has interior completely inside the Teichmüler space, moreover, by the local compactness of the WP metric at x after possibly passing to a subsequence the initial parts of r_n limit to a ray r starting at x.

We would like to show that r is an infinite ray. Let $s_n = \frac{1}{\ell_{\rho(n)}(x)}$. Then for each n and t in the domain of r_n we have $\ell_{s_n\rho(n)}(r_n(t)) \leq 1$. This follows from the convexity of length-functions along WP geodesics and the observation that $\ell_{s_n\rho(n)}(x) = 1$ and $\ell_{s_n\rho(n)}(c_n) = 0$. Now assuming that $r(t)$ has finite length T we get a contradiction.

Assume that after possibly passing to a subsequence $s_n\rho(n) \to \mu$ in the weak* topology of $\mathcal{ML}(S)$. We claim that $|\mu|$ fills Z. To see this note that for n sufficiently large $\rho(n) = \partial Z \cup v_n$ where $v_n \in g_Z$ the infinite geodesic of ρ, then by Theorem 2.2 $s_n\tau_n$ after possibly passing to a subsequence limits in the weak* topology of $\mathcal{ML}(Z)$ to a measured lamination μ' whose support by is equivalent to ν'. Thus $|\mu| \subset |\nu'| \cup \{$ some curves in $\partial Z\}$ and the claim follows.

The claim implies that every γ with $\iota(\gamma, \mu) = 0$ is in ∂Z (recall that Z is non-separated).

Moreover, the length-functions $\ell_\mu(x) : \text{Teich}(S) \times \mathcal{ML}(S) \to \mathbb{R}^{\geq 0}$ are continuous in both x and μ, which implies that for any $t < T$ we have

$$\ell_\mu(r(t)) \leq 1 \tag{7.1}$$

Let σ be the simplex in $\mathcal{C}(S)$ such that $r(T) \in \mathcal{S}(\sigma)$.

We note that $\lim_{t\to T} \ell_\mu(r(t)) \leq 1$ which implies that for each simple closed curve $\gamma \in \sigma$, we have $\iota(\gamma, \mu) = 0$. Otherwise since $\ell_\gamma(r(t)) \to 0$ as $t \to T$, we would have $\ell_\mu(r(t)) \to \infty$ which contradict the bound (7.1). Therefore,

$$\sigma \subseteq \partial Z$$

Note that $r_n(T) \to r(T)$ as $n \to \infty$. Now we verify the following two claims:

Claim 1: The points c_n for n sufficiently large lie in $\overline{\mathcal{S}(\sigma)}$. To see this note that for n sufficiently large $\partial Z \subset \rho(n)$, also as we saw above $\sigma \subseteq \partial Z$, thus $\sigma \subset \rho(n)$ for n sufficiently large. Moreover, c_n is maximally pinched at $\rho(n)$, and the claim follows.

Claim 2: The distance from $r_n(T)$ to c_n in the completion is diverging. By Brock's quasi-isometry we only need to show that $d(Q(r_n(T)), \rho(n)) \to \infty$. To see this note that by the $D = D(A)$ fellow-traveling, Theorem 5.15, and Brock's quasi-isometry every $Q(r_n(T))$ is in distance at most $K_{\text{WP}}T + C_{\text{WP}} + D$ from $\rho(0)$, moreover $d(\rho(0), \rho(n)) \to \infty$ as $n \to \infty$.

Now we claim that for any $t' > T$ the points $r_n(t')$ converge into the $\sigma-$stratum. To see this note that we have $r_n(T) \to r(T) \in \mathcal{S}(\sigma)$ as $n \to \infty$, by claim 1 above and since $\mathcal{S}(\sigma)$ is geodesically convex for any n sufficiently large $[r(T), c_n] \subset \mathcal{S}(\sigma)$, by claim 2 $d_{\mathrm{WP}}(r_n(T), c_n) \to \infty$. Now since $r_n(t')$ are points with fixed distance $t' - T$ from $r_n(T)$ the claim follows from a CAT(0) comparison for the triangles $\triangle(r_n(T)r(T)c_n)$.

Therefore the finite-length geodesic segments $r_n([0,t'])$ have endpoints converging in the completion, and thus these limiting endpoints are the end points of a geodesic segment which by the non-refraction theorem its interior lies in the maximal stratum containing its endpoints.

Since one endpoint of each geodesic $r_n([0,t'])$ is the base surface x, the interior of the limit segment lies in the interior of Teichmüller space. Moreover, the approximating geodesics $r_n([0,t'])$ which are parametrized by arc-length converge to the parametrization by arc-length of the limit segment. Thus, the limit $r(T)$ of the points $r_n(T)$ lies in the interior of this geodesic limit, contradicting the assumption that $r(T) \in \mathcal{S}(\sigma)$. This finishes the proof of that the geodesic r is an infinite geodesic ray.

□

In the following theorem we show that the behavior of r_ρ mimics combinatorial properties of hierarchy paths encoded in the ending data. We say that a subsurface Z has $(R, R')-$bounded combinatorics over a subinterval of r_ρ, if r_ρ fellow-travels ρ over a subinterval where the subsurface Z has $(R, R')-$bounded combinatorics.

Theorem 7.2. *(Infinite ray with prescribed itinerary) Let $\rho : [0, \infty) \to P(S)$ be an infinite hierarchy path with $A-$narrow ending data, and $r_\rho : [0, \infty) \to$ Teich(S) be the corresponding infinite WP geodesic ray constructed in Lemma 7.1.*

Given positive constants R and R' and $\epsilon > 0$ sufficiently small there are constants $\bar{s}$ and $\bar{\epsilon}$ such that if a non-separated subsurface Z has $(R, R')-$bounded combinatorics over an interval of length $2\bar{s} + a$ for any positive a then there is an interval $[t^-, t^+]$ of length at least a such that

1. *$\ell_{\partial Z}(r_\rho(t)) \leq \epsilon$ and*
2. *$\ell_\gamma(r_\rho(t)) > \bar{\epsilon}$ for every $\gamma \notin \partial Z$*

for every $t \in [t^-, t^+]$.

Moreover, if there are several domains Z as above then the time intervals show up along r_ρ in the same order as the order of intervals along the hierarchy path ρ.

Proof. For each n the ending data of the geodesic segment $r_n = [x, c_n]$ is $(\rho(0), \rho(n))$ which by Theorem 2.11 (5) is $A + 2M_2$ narrow, thus Theorem 6.1 (1) gives the upper bound in (1), and Lemma 6.8 (2) gives the lower bound (2), note that $s \ll \bar{s}$ and the lower bound holds on a much longer interval.

Now the continuity of length-functions on Teichmüller space implies that along the limit geodesic ray r_ρ we have the same bounds on length-functions. Also the intervals show up in the order in which subsurfaces show up along ρ. We will refer to this ray as a *ray with prescribed itinerary.* $\square$

7.2 Diverging geodesic rays

In this subsection we construct diverging Weil-Petersson geodesics in the moduli space. A ray is diverging if eventually leaves every compact subset of moduli space. More precisely

Definition 7.3. (Diverging ray) Let $r : [0, \infty) \to \mathcal{M}(S)$ be a ray in the moduli space, then r diverges if for every $\epsilon > 0$ there is $T \in [0, \infty)$ such that $r([T, \infty))$ be in the $\epsilon-$thin path of moduli space.

Theorem 1.2 in introduction which asserts that there are uncountably many diverging rays with minimal filling ending lamination is a consequence of our construction in this subsection.

We use the results and notations from appendix. Let $\alpha_1, \alpha_2 \in \mathcal{C}(S)$ be disjoint, moreover α_1, α_2 and $\alpha_1 \cup \alpha_2$ be non-separating on S. Let $X_1 = S - \alpha_1$, $X_2 = S - \{\alpha_1, \alpha_2\}$ and $X_3 = S - \alpha_2$. Moreover as in the appendix let each f_i for $i = 1, 2, 3$ be a pseudo-Anosov map supported on X_i.

The constructions of this section are quite general and apply to any collection of non-separated subsurfaces which intersect each other in a specific pattern (see Figure 7), we restrict to the special case to give a simpler and more clear discussion.

Let $\bar{\tau} = \min_{1=1,2,3} \tau_{X_i}(f_i)$ be as in Proposition 9.4 and m be as in Proposition 9.7.

Consider the function $q : \mathbb{N} \to \{1, 2, 3\}$ where $q(i) \equiv i \pmod 3$ and any $e : \mathbb{N} \to \mathbb{Z}$.

Then by Proposition 9.7 all coefficients of the ending data $\nu^- = \mu^-_{q,e}$ and $\nu^+ = \mu^+_{q,e}$ assigned to separated subsurfaces are bounded by m, now since there is no separated subsurface is in the list $\{X^{q,e}(f_{q(i)})\}_i$ then the ending data is $m-$narrow. Let K and C be the constant from Proposition 5.16.

Let r be the WP geodesic ray with same ending data and prescribed itinerary.

Let $\epsilon_i \to 0$ be a decreasing sequence going to 0 and $e_i \to \infty$ be an increasing sequence of integers going to ∞ such that

$$\bar{\tau} e_i > K \bar{s}(m, m, m, \epsilon_i) + C$$

where $\bar{s}$ is the constant from Theorem 7.2, note that for fixed m,

$$\bar{s}(m, m, m, \epsilon_i) \to \infty \text{ as } \epsilon_i \to 0$$

Then as we saw in the appendix along the infinite hierarchy given by the function q and the sequence $\{e_i\}_i$ as above, each $Z_i = X^{q,e}(f_{q(i)})$ has $M-$bounded combinatorics over an interval of length at least $\bar{s}(m, m, m, \epsilon_i)$.



Figure 7: Illustration of a more general intersection pattern of component domains a hierarchy path ρ with diverging corresponding ray r_ρ: Over each red subinterval a non-separated domain has bounded combinatorics. For each i, $J_{Z_i} \cap J_{Z_{i+2}} \neq \emptyset$ because a non-separated subsurface Z_{i+1} of $Z_i \cap Z_{i+2}$ is a component domain, Z_{i+1} has bounded combinatorics over the corresponding red subinterval. The red subintervals get longer and longer forward in time.

Now Theorem 7.2 implies that there are times $t_1 < ... < t_i < t_{i+1} < ...$ such that

$$\ell_{\partial Z_i}(g(t_i)) < \epsilon_i$$

Moreover, it is easy to check that any two consecutive domains Z_i and Z_{i+1} have a boundary curve in common, so by convexity of length-functions

$$\ell_{\partial Z_i}(r(t)) < \epsilon_i \text{ for all } t \in [t_i, t_{i+1}]$$

Now since the intervals $[t_i, t_{i+1}]$ cover $[0, \infty)$ and $\epsilon_i \to 0$ the WP geodesic ray diverges in Teichmüller space which implies that its projection to the moduli space also diverges.

Further the forward ending lamination of r_ρ is minimal filling because the main geodesic of ρ is an infinite geodesic ray in $\mathcal{C}(S)$, which determine a minimal filling lamination in the Gromov boundary of $\mathcal{C}(S)$.

7.3 Closed geodesics in the thin part

In this subsection we provide examples of closed Weil-Petersson geodesics which stay in the thin part of moduli space. Theorem 1.1 in the introduction follows from our construction, because any compact subset of moduli space $\mathcal{K}$ is contained in the $\epsilon = \inf_{x \in \mathcal{K}} \mathrm{inj}(x)$ thick part of moduli space.

Let curves α_1 and α_2, sub-surfaces X_1, X_2 and X_3 and pseudo-Anosov maps f_1, f_2 and f_3 be as in the previous subsection. Let function $q : \mathbb{N} \to \{1, 2, 3\}$ be defined as $q(i) \equiv i \pmod 3$, moreover let $e : \mathbb{N} \to \mathbb{N}$ be such that $e_i = e_{i+3}$ and for $i = 1, 2, 3$ e_i be large enough such that

$$\bar{\tau} e_i > K \bar{s}(m, m, m, \epsilon) + C$$

then the corresponding WP geodesic stays in the $\epsilon-$thin part of moduli space. Therefore its projection to the moduli space which is a closed WP geodesic stays in the $\epsilon-$thin part of moduli space.

7.4 A recurrence criterion

The following criterion is a straightforward consequence of Corollary 6.10 and Theorem 7.2. Given constants A, R and R' recall the constants $s = s(A, R)$ and $\bar{\epsilon} = \bar{\epsilon}(A, R, R')$ form the corollary and let $L = Ks + C$ where K and C are the constants from Proposition 5.16.

Theorem 7.4. *(Recurrence cirterion) Let ρ be an $A-$narrow hierarchy path, let I_k be a sequence of intervals of length at least L over each of which S has $(R, R')-$bounded combinatorics. Then the ray with prescribed itinerary and the same ending data is recurrent to the $\bar{\epsilon}-$thick part of moduli space.*

Here we give a construction for which this criterion applies.

Let $\{X_i\}_{i=1}^n$ be a collection of non-separated subsurfaces of S and $X_1 = S$, pseudo-Anosov maps $\{f_i\}_{i=1}^n$ such that f_i is supported on X_i. Let $q : \mathbb{N} \to \{1, 2, ..., n\}$, also constants M and $\bar{\tau} = \min_{i=1,\dots,n} \tau(f_i)$ be as in the appendix. Assume that for a sequence of integers $\{i_k\}_k$, $q(i_k) = 1$ then $f_{q(i_k)}$ is supported on S. Now by the above theorem and if e_{i_k} be large enough such that

$$\bar{\tau} e_{i_k} > Ks(m, m, m) + C$$

then the geodesic ray corresponding to q and e is recurrent to the $\bar{\epsilon}(m, m, m)$ thick part of moduli space.

Note that if $X_2 \subsetneq S$ be a proper subsurface S and for another subsequence j_k $q(j_k) = 2$, then if $e_{j_k} \to \infty$ the ray corresponding to q and e is recurrent but no compact subset of moduli space contains this ray.

8 Asymptotic class of rays with narrow ending data

In this section we use the ruled surface arguments similar to [BMM10] to prove Theorem 1.3 which is the ending lamination theorem for WP geodesics with narrow ending data and bounded annular coefficients,

Theorem 1.3. *The strong asymptotic class of a WP geodesic ray with narrow ending data and bounded annular coefficients is determined by the forward ending lamination ν^+.*

This theorem is a generalization of Brock, Masur and Minsky's *Recurrent ending lamination theorem* in [BMM10], asserting that the strong asymptotic class of recurrent WP geodesic rays to a compact subset of moduli space is determined by the forward ending lamination. These ending lamination theorems are partial results toward a parametrization of the visual boundary of the WP metric in terms of laminations.

8.1 Bounding annular coefficients

Let r_ρ be a WP geodesic ray with prescribed itinerary with narrow ending data with bounded annular coefficients, we show that given $a > 0$ and $\epsilon > 0$ sufficiently small there is a subinterval of length at least a of any long enough subinterval of r_ρ over which r_ρ visits some $U_{\epsilon,\bar{\epsilon}}$ region. By Theorem 7.2 we only need to find long enough subintervals of the hierarchy path ρ where a domain Z has bounded combinatorics.

Lemma 8.1. *Given an increasing function $F : \mathbb{R}^{\geq 0} \to \mathbb{R}^{\geq 0}$ there is a constant L, depending only on F and the topological type of S with the following property:*

Let $\rho : [m, n] \to P(S)$ be a hierarchy path, let a subinterval $[m', n'] \subseteq [m, n]$ has the property that for any subsurface Z and any $R > 0$ any subinterval of $[m', n']$ where Z has non-annular $R-$bounded combinatorics has length at most $F(R)$. Then

$$n' - m' < L$$

Proof. For each $1 \leq i \leq \xi(S)$ define

- $x_i = \max\{d_Y(\rho(m'), \rho(n')) : Y \subseteq S$ is non-annular and $\xi(Y) \leq i\}$

For each $1 \leq i \leq \xi(S)$ if there is a non-annular subsurface Z with $\xi(Z) = i$ such that $d_Z(\rho(m'), \rho(n')) > F(x_{i-1}) + 2M_2$ then by Theorem 2.11 (4),(5) $|J_Z \cap [m', n']| > F(x_{i-1})$. Thus Z has non-annular $x_{i-1}-$bounded combinatorics over the subinterval $J_Z \cap [m', n']$ of length at least $F(x_{i-1})$ which is not possible by the assumption of lemma. Thus for every Z with $\xi(Z) = i$ we have $d_Z(\rho(m'), \rho(n')) \leq F(x_{i-1}) + 2M_2$, now since

$$x_i = \max\{x_{i-1}, \max_{Z:\xi(Z)=i} d_Z(\rho(m'), \rho(n'))\}$$

we get the following bounds:

$$x_i \leq \max\{x_{i-1}, F(x_{i-1}) + 2M_2\} \text{ for every } 1 \leq i \leq \xi(S)$$

for simplicity we let $f(x) = \max\{x, F(x) + 2M_2\}$, then using the above inequalities inductively we get

$$x_i \leq f^{(i)}(0) \tag{8.1}$$

for every $1 \leq i \leq \xi(S)$

Moreover, from the layered structure of hierarchies and Theorem 2.11 (5) we get

$$n' - m' \leq \prod_{i=0}^{\xi(S)} (x_i + 2M_2)$$

(see [Min01] Theorem 5.1 for a discussion about the above bound). Thus by the bounds (8.1) we get

$$n' - m' \leq \prod_{i=0}^{\xi(S)} (f^{(i)}(0) + 2M_2)$$

which is the desired bound on the length of $[m', n']$ only depending on F and the topological type of S.

□

Lemma 8.2. *Given positive constants A, a, R' and $\epsilon \leq \bar{\epsilon}$, there is $L' > 0$ with the following property:*

Let $\rho : [0, \infty) \to P(S)$ be a hierarchy path with $A-$narrow ending data with $R'-$bounded annular coefficients. Let $r_\rho : [0, \infty) \to \mathrm{Teich}(S)$ be the ray prescribed with the same ending data, then for any subinterval of length at least L' there is a non-separated domain Z such that r_ρ visits $U_{\epsilon,\bar{\epsilon}}(\partial Z)$ over an interval $I = [t^-, t^+]$ of length at least a.

Proof. Let K and C be the constants from Proposition 5.16. Define

$$F(x) = \max\{K\big(2\bar{s}(A, x, R', \epsilon) + a\big) + C, A\}$$

where $\bar{s}$ is the function from Theorem 1.5.

Take a subinterval of the ray of length at least $L' = KL + C$, then Lemma 8.1 implies that there are $R > 0$, subsurface Z and an interval $I' \subset [0, \infty)$ with $|I'| \geq F(R)$ such that Z has non-annular $R-$bounded combinatorics over I'. Moreover, since ρ is $A-$narrow by the definition of F the subsurface Z is non-separated.

By the assumption of the lemma we have the bound R' on the annular coefficients over $[m', n']$. Therefore, Z has $(R, R')-$bounded combinatorics over $[m', n']$, thus the lemma follows form the bounds on length-functions in Theorem 7.2. □

In the following corollary we consider the case that ρ has an infinite geodesic in a proper subsurface $Z \subsetneq S$.

Corollary 8.3. *Let $\rho : [0, \infty) \to P(S)$ be a half-infinite hierarchy path as in the previous lemma with an infinite tight geodesic in $\mathcal{C}(Z)$ ($Z \subsetneq S$). Let r_ρ be a WP geodesic ray . Then*

$$\ell_{\partial Z}(r(t)) \to 0$$

as $t \to \infty$.

Proof. By the previous lemma if we let $\epsilon_k \to 0$ for longer and longer subintervals along ρ there are subsurface Z_k such that $\inf_{t \in I_k} \ell_{\partial Z_k} r_\rho(t) \leq \epsilon_k \to 0$. Moreover by the $A-$narrow condition each subsurface Z_k is non-separated, also by the structure of hierarchy paths for every k sufficiently large $\partial Z \subseteq \partial Z_k$. Thus the the corollary follows from the convexity of length-functions. □

8.2 Strongly asymptotic rays

In this subsection we continue to prove Theorem 1.3. Our strategy is the same as [BMM10]. In the *narrow visibility theorem*, Theorem 8.9, we show that given a WP geodesic ray r_ρ with prescribed itinerary with narrow ending data with bounded annular coefficients and r' a geodesic with the same forward ending data and not strongly asymptotic to r_ρ there is a bi-infinite geodesic g asymptotic to r_ρ in the forward time and to r' in backward time. In Theorem 8.7 we show that g is actually strongly asymptotic to r_ρ. These together will lead to a contradiction to the convexity of the length of measured laminations along the WP geodesic g, so we conclude that r_ρ and r' are strongly asymptotic and the forward ending data determines the strong asymptotic class of ray with narrow ending data and bounded annular coefficients.

The main new ingredient here is a kind of geometric limit argument which we use to produce infinitely many regions with a definite total curvature marching off to infinity along r_ρ on ruled surfaces consisting of smooth families of geodesic segments orthogonal to r_ρ. In [BMM11] the recurrence to the thick part was used to produce such regions. Although the sectional curvatures of the WP metric are not bounded away from 0 this technique helps us to prove visibility of this class of rays, for some standard visibility arguments see [Ebe72].

Definition 8.4. Two geodesic rays $r, r' : [0, \infty) \to \mathcal{X}$ in a metric space parametrized by arc-length are asymptotic if there are d and T positive such that for every $t \geq T$:

$$d(r(t), r'(t)) \leq d$$

r and r' are strongly asymptotic if

$$d(r(t), r'(t)) \to 0$$

as $t \to \infty$

Here we present some results about ruled surfaces in negatively curved spaces.

Let $r : I \to \mathcal{X}$ be a geodesic in a complete non positively curved Riemannian manifold or orbifold (for example moduli space equipped with WP metric), let $\mathcal{R}$ be a ruled rectangle obtained from a smooth family of geodesic segments of length at least δ orthogonal to $r(I)$.

For every $s \in [0, \delta]$ let l_s be the length of the path of equidistance s from $r(I)$ in $\mathcal{R}$, and let $u_s : [0, 1] \to \mathcal{R}$ be parametrization of this path by constant speed l_s.

Let $u : [0, 1] \times [0, \delta] \to \mathcal{X}$ be the parametrization of $\mathcal{R}$ defined by $u(t, s) = u_s(t)$. For every $t \in [0, 1]$ we denote the path $u(t, .)$ by $u^t : [0, \delta] \to \mathcal{R}$.

The family of paths u^t and u_s consist respectively a vertical and a horizontal foliation of $\mathcal{R}$ which are orthogonal to each other, see Figure 8.

Let $p_s : u_s([0, 1]) \to u_0([0, 1])$ be the projection obtained by following the ruling geodesic paths u^t.

For every $s \in [0,\delta]$ denote by $\mu_{\mathcal{R},s}$ the measure obtained by integrating the function $k_g(u_s(t))$ on u_s with respect to da_s the arc-length. Note that since each u_s is concave in $\mathcal{R}$ the function $k_g(u(s))$ is a positive.

Define $\mu_{\mathcal{R}} = \lim_{s\to 0} p_{s*}(\mu_{\mathcal{R},s})$ where p_{s*} is the push forward defined by p_s from the Lebesgue measures on $u_s([0,1])$ to the ones on $u_0([0,1])$.

Given a subinterval $I' \subseteq [0,1]$ by the Gauss-Bonnet theorem (see [Kli78]) for $u(I' \times [s,\delta])$ we have

$$\int\int_{u(I'\times[s,\delta])} \kappa dA + \int_{\partial u(I'\times[s,\delta])} k_g da + \sum_i \theta_i = 2\pi$$

where the sum is over the exterior angles at the four corners of $\partial u(I' \times [s,\delta])$. By the orthogonality of foliations the exterior angles are each $\frac{\pi}{4}$ which add up to 2π. Moreover, the sides $u^0([0,\delta])$ and $u^a([0,\delta])$ are geodesics in $\mathcal{X}$ and consequently the the intrinsic metric of $\mathcal{R}$, thus we get

$$\int\int_{u(I'\times[s,\delta])} +\kappa dA - \int_{u_s(I')} k_g da_s = \int_{u_0(I')} k_g da_0$$

Now taking limit as from the above equality $\epsilon \to 0$ we get

$$\int_{u_0(I')} \mu_{\mathcal{R}} = \int\int_{u(I'\times[s,\delta])} -\kappa dA + \lim_{s\to 0}\int_{u_s} k_g da_s$$

but $\lim_{s\to 0}\int_{u_s(I')} k_g da_s = \int_{u_0(I')} \mu_{\mathcal{R}}$, thus we have

$$\int_{r(I')} \mu_{\mathcal{R}} = \int\int_{u(I'\times[s,\delta])} -\kappa dA + \int_{u_0(I')} \mu_{\mathcal{R}} \tag{8.2}$$

By the CAT(0) comparison for the rectangle $u([t',t]\times[0,s])$ the distance between any two geodesic segments $u^{t'}(s)$ and $u^t(s)$ is increasing in s, thus $l_s \geq l_{s'}$ for $s' \leq s$.

Lemma 8.5. *For any $I' \subseteq I$ and $s \in [0,\delta]$ let*

$$\phi(s) = \int\int_{u(I'\times[0,s])} -\kappa dA + \int_{r(I')} \mu_{\mathcal{R}}$$

Then ϕ is an increasing function in s and

$$\phi(s) \geq \frac{l_s - l_0}{s}$$

Proof. The fact that ϕ is an increasing function in s follows from $-\kappa \geq 0$.

Here we prove the following integral formula:



Figure 8: The ruled rectangle $\mathcal{R}$, the vertical path are the equidistant paths from $r(I)$ and horizontal paths are the ruling geodesic segments which consist a pair of orthogonal foliations for $\mathcal{R}$.

$$\begin{aligned} l_s - l_0 &= \int_0^s \frac{dl_s}{ds} ds \qquad (8.3) \\ &= \int_0^s \Big(\int_{u_s} k_g(u_s(a)) da_s \Big) ds \\ &= \int_0^s \Big(\int\int_{u(I' \times [0,s])} -\kappa dA + \int_{u_0(I')} \mu_{\mathcal{R}} \Big) ds \end{aligned}$$

The first equality is the fundamental theorem of calculus applied to l_s as a function of s.

The second equality holds by the first variation of arc-length formula for an orthogonal variation of paths, see for example [CE08] page 4, where $k_g(u_s(t))$ is the geodesic curvature of the path u_s at the point $u_s(t)$ in the ruled surface $\mathcal{R}$. Here da_s is the arc-length element along u_s.

The last equality holds by (8.2).

Now by the formula (8.3) and since $\phi(s)$ is an increasing function for every $s \in [0, \delta]$ we have

$$l_s - l_0 \leq \phi(s)s \qquad (8.4)$$

Therefore, $\phi(s) \geq \frac{l_s - l_0}{s}$ and the lemma follows. □

Remark 8.6. If the ruling extends to a ruled surface containing $r(I)$ in its interior then the second term in $\phi(s)$ vanishes. Defining the measure $\mu_{\mathcal{R}}$ and the rather long argument above meant to handle the possibility that the ruling does not extend which could be the case in the following theorems.

Now we continue to prove our results in the setting of the WP metric.

Theorem 8.7. *(Asymptotic $\Longrightarrow$ strongly asymptotic)*

Let r_ρ be a WP geodesic ray with prescribed itinerary with $A-$narrow ending data and bounded annular coefficients. Then any WP geodesic ray r' asymptotic to r_ρ is also strongly asymptotic to r_ρ.

Proof. For simplicity of notation we denote r_ρ by r. Assume that $r(t)$ and $r'(t)$ are parametrized by arc-length.

We may assume that there is $\delta > 0$ such that the distance between $r(t)$ and $r'(t)$ is bounded below by δ, otherwise r' and r would be strongly asymptotic.

Lemma 8.2 implies that given $A, a > 0$ and a sequence $\epsilon_k \to 0$, there are time intervals $I_k = [t_k^-, t_k^+]$ of length a (we can always assume that the intervals are of length a because a priori they have length at least a) and non-separated subsurfaces Z_k such that $r(I_k) \subseteq U_{\epsilon_k,\bar{\epsilon}}(\partial Z_k)$, moreover $t_k^+ < t_{k'}^-$.

Now consider the ruled surface $\mathfrak{R}_n \subset \mathrm{Teich}(S)$ as follows: $\mathfrak{R}_n$ consists of a smooth family of geodesic segments orthogonal to the geodesic segment $(r(0)r(n))$ from $(r(0)r'(n)) \cup (r'(n)r(n))$, see Figure 9.

In what follows we set up a technique to produce a family of ruled rectangles on the ruled surfaces $\mathfrak{R}_n$ with the property that all of them have a definite negative total curvature. These rectangles have one side $r(I_k)$ and march to infinity along r.

We denote the projections of r and r' to moduli space, respectively by r and r' as well.

First note that $\overline{\mathcal{M}(S)}$ is compact and each $r|_{I_k} : [0, a] \to \overline{\mathcal{M}(S)}$ is $1-$Lipschitz (each one is a geodesic segment), thus $r|_{I_k}$ is an equicontinuous family of maps, and therefore by the Arzelá-Ascoli theorem (see [Mun75]) after possibly passing to a subsequence converge point-wise as parametrized geodesics to a geodesic segment $\bar{r} : [0, a] \to \overline{\mathcal{M}(S)}$. Lifting the picture to Teich(S) since $r(I_k) \subset U_{\epsilon_k,\bar{\epsilon}}(\partial Z_k)$ for non-separating multi-curves ∂Z_k and $\epsilon_k \to 0$, thus $\bar{r}([0, a]) \subset \mathcal{S}(\sigma)$ for a non-separating multi-curve σ.

Since the rays r and r' are not strongly asymptotic we may omit finitely many of I_k and assume that there is $\delta > 0$ such that for each k the distance of the geodesic segments $r(I_k)$ from r' is at least δ. To see this note that in a CAT(0) space the distance between any two geodesic rays is a monotonic function.

Now for every n there is a collection of ruled rectangle $\mathcal{R}_{k,n}$ in $\mathfrak{R}_n$ consisting of the terminal subsegments of the ruling geodesic segments of $\mathfrak{R}_n$ of length δ with end point on $r(I_k)$, let $u_{k,n} : [0, 1] \times [0, \delta] \to \mathfrak{R}_n$ be the parametrization of $\mathcal{R}_{k,n}$ as in the discussion before this theorem, see Figure 9. Moreover, for every k and n as above define

$$\phi_{k,n} = \int\int_{\mathcal{R}_{k,n}} -\kappa dA + \int_{r(I_k)} \mu_{\mathcal{R}_{k,n}}$$

We continue to show that for some $K_0 > 0$

$$\phi_{k,n} \geq K_0 \tag{8.5}$$



Figure 9: Building up negative curvature on the ruled surfaces $\mathfrak{R}_n$. The bold subintervals $r(I_k) \subset U_{\epsilon,\bar{\epsilon}}(\partial Z_k)$ where Z_k is a non-separated component domain of the hierarchy path ρ. Ruled rectangle $\mathcal{R}_{k,n}$ consists of subsegments of length δ of the ruling geodesic segments of $\mathfrak{R}_n$ terminating on $r(I_k)$.

holds.

The proof is by contradiction. First assume that for some $s \in [0, \delta]$ and a sequence of pair of integers (k_i, n_i)

$$(*)\phi_{k_i,n_i} \to 0 \text{ as } i \to \infty$$

To simplify notation in the rest of the proof we denote u_{k_i,n_i} by u_i, ϕ_{k_i,n_i} by ϕ_i, also for any $s \in [0, \delta]$ denote $u_{k_i,n_i,s}$ by $u_{i,s}$ and $l_{k_i,n_i,s}$ by $l_{i,s}$.

First assume that for some $s \in [0, \delta]$ $u_{i,s}$ is not equicontinuous, then there is $\epsilon > 0$, a sequence $\delta_m \to 0$ and a sequence of integers i_m such that

$$d_{\mathrm{WP}}(u_{i_m,s}(t + \delta_m), u_{i_m,s}(t)) > \epsilon$$

moreover, $l_{i_m,s}\delta_m \geq d_{\mathrm{WP}}(u_{i_m,s}(t + \delta_m), u_{i_m,s}(t))$. Now for every m sufficiently large $\frac{\epsilon}{\delta_m} - a > s$ thus Lemma 8.5 for the ruled rectangle $u_i([0, 1] \times [0, s])$ implies that

$$\int\int_{u_i([0,1]\times[0,s])} -\kappa dA + \int_{u_i([0,1])} \mu_{\mathcal{R}_i} > 1$$

moreover ϕ_i is larger than the left hand side of the above inequality, so we get the contradiction to (*).

Thus in the rest of proof we can assume that for every $s \in [0, \delta]$ $u_{i,s}$ is an equicontinuous family of maps, which implies that $u_i : [0, 1] \times [0, \delta] \to \overline{\mathcal{M}(S)}$ is equicontinuous. Therefore by the Arzelá-Ascoli theorem after possibly passing to a subsequence u_i converge point-wise to a map $\bar{u} : [0, 1] \times [0, \delta] \to \overline{\mathcal{M}(S)}$. For definition of an equicontinuous family of maps and the statement of the Arzelá-Ascoli theorem see [Mun75].

Recall that $r|_{I_k}$ limit into the $\epsilon-$thick part of $\mathcal{S}(\sigma)$ so $u_{i,0}|_{[0,1]}$ which is a unit speed reparametrization of $r|_{I_k}$ limit into the $\epsilon-$thick part of $\mathcal{S}(\sigma)$ as well.

By Lemma 3.5 there is a lower bound for the distance the of $\bar{u}|_{[0,1]}$ and any stratum $\mathcal{S}(\tau)$ where $\tau \not\subseteq \sigma$, thus there is $\epsilon' < \epsilon$, simplex $\sigma' \subseteq \sigma$ and intervals $I_1 \subset [0,1]$ and $I_2 \subset [0,\delta]$ such that $\bar{u}(I_1 \times I_2)$ lies in the $\epsilon'-$thick part of $\mathcal{S}(\sigma')$, see Figure 10.

$\mathcal{S}(\sigma')$ is totally geodesic thus by continuity of distances in a point-wise converging sequence $\bar{u}_s|_{[0,1]}$ is the path of equidistance s from the path $\bar{u}_0|_{[0,1]}$ in $\mathcal{S}(\sigma')$, we denote the length of $\bar{u}_s|_{[0,1]}$ by $\bar{l}_s$.

Moreover since σ' is non-separating the sectional curvatures are all negative and bounded away from 0 by some $\kappa' < 0$ in the $\epsilon'-$thick part of $\mathcal{S}(\sigma')$, see [Wol03].

Now using the comparison with CAT(κ') space we have

$$\bar{l}_s - a > as(-\kappa')$$

To see the above inequality, let $\bar{l}_s$ be the length of paths in an equidistant family as $\bar{u}_s$ in a CAT(κ') space with $\kappa' < 0$. Using the Fermi coordinates (see [Kli78]) one can see that $\bar{l}_s = \sinh(-\kappa' d)\bar{l}_0$, then the above inequality follows since $\sinh x > x$ for every $x > 0$.

Now since $u_{i,s}$ converge point-wise to u_s, $l_{i,s} \to \bar{l}_s$ as $i \to \infty$, so for every i sufficiently large $l_{i,s} - l_{0,s} = l_{i,s} - a > \frac{-as\kappa'}{2}$, so Lemma 8.5 for the ruled rectangle $u_i(I_1 \times I_2)$ implies that

$$\int\int_{u_i(I_1 \times I_2)} -\kappa dA \geq \frac{-as\kappa'}{2s} = \frac{-a\kappa'}{2}$$

moreover ϕ_i is greater than the left hand side of the above inequality, so we get the contradiction to (*).

So we are done with proving that the bound in (8.6) holds for all $\mathcal{R}_{k,n}$.

Remark 8.8. In the case that $\sigma' = \emptyset$, we can get the contradiction alternatively as follows: for every k sufficiently large $q_k = u_k^t(s)$ is in some $\epsilon' < \epsilon$ thick part of Teichmüller space. Thus there are $r > 0$ and $\kappa' < 0$ such that for every k sufficiently large the curvature on $\mathcal{N}_r(q_k) \cap \mathcal{R}_{k,n}$ is bounded by κ'. Moreover, the area of $\mathcal{N}_r(q_k) \cap \mathcal{R}_{k,n}$ is at least $\pi\frac{r^2}{2}$ so $\int\int_{\mathcal{N}_r(q_k)\cap\mathcal{R}_{k,n}} -\kappa dA > \frac{\pi r^2}{2}(-\kappa')$, thus again since ϕ_i is larger than the left hand side we get the contradiction to (*).

We continue to show that r and r' are strongly asymptotic. Assume they are not strongly asymptotic then for some $\delta_0 > 0$ and T

$$d_{\mathrm{WP}}(r(t), r'(t)) > \delta_0$$

for all $t \geq T$. Moreover, since r and r' are asymptotic there is d such that $d_{\mathrm{WP}}(r(t), r'(t)) \leq d$. Let $\delta = \frac{\delta_0}{2}$ then the CAT(0) comparison for the ruled triangle $\mathfrak{R}_n$ shows that for n sufficiently large N the number of ruled rectangles $\mathcal{R}_{k,n}$ with hight δ as above such that $\mathcal{R}_{k,n} \subset \mathfrak{R}_n$ would be made arbitrary large.

Further by (8.6) we have the estimate



Figure 10: σ is a non-separating multi-curve and $u_{i,0}|_{[0,1]}$ limit into the $\epsilon-$thick part of $\mathcal{S}(\sigma)$. $\sigma' \subseteq \sigma$ and $u_i(I_1 \times I_2)$ limit to the smaller ruled rectangule in the $\epsilon'-$thick part of $\mathcal{S}(\sigma')$. Moreover, each $u_{i,s}|_{I_1}$ limits to a path of equidistance s from $u_{i,0}|_{I_1}$

$$\int\int_{\mathfrak{R}_n} -\kappa dA + \int_{r([0,n])} \mu_{\mathfrak{R}_n} > NK_0$$

Here as in the discussion before this theorem $\mu_{\mathfrak{R}}$ is defined as the limit of push forwards of measures obtained by integrating geodesic curvature of equidistant paths in $\mathfrak{R}_n$ along the ruling geodesics in $\mathfrak{R}_n$.

Moreover, by the Gauss-Bonnet theorem the left hand side is bounded above by π independent of N, so we get the upper bound $\frac{\pi}{K_0}$ for N.

This contradicts the observation above that N would be made arbitrary large.

Now since $\delta_0 > 0$ was arbitrary we conclude that the rays r and r' are strongly asymptotic. $\square$

Theorem 8.9. *(Narrow visibility)*

Let r_ρ be a WP geodesic ray with prescribed itinerary with narrow ending data and bounded annular coefficients, let r' be a WP geodesic ray with the same forward ending lamination which is not strongly asymptotic to r. Then there is a bi-infinite geodesic inside Teichmüller space which is strongly asymptotic to r_ρ in forward time and asymptotic to r' in backward time.

Proof. Again for simplicity we denote r_ρ by r. Let r and r' both be parametrized by arc-length. Since the rays r and r' are not strongly asymptotic we may omit finitely many of I_k and assume that there is $\delta > 0$ such that for each k the distance of the geodesic segments $r(I_k)$ from r' is at least δ. To see this note that in a CAT(0) space the distance between any two geodesic rays is a monotonic function.

For any integer $n > 0$ let the rued surfaces $\mathfrak{R}_n$ be as in the previous theorem and denote by g_n the parametrization by arc-length of $(r(n)r'(n))$. We define $u_{k,n} : [0,1] \times [0,\delta] \to \overline{\mathcal{M}(S)}$ and $\mathcal{R}_{k,n} = u_{k,n}([0,1] \times [0,\delta]) \subset \mathfrak{R}_n$ as in the previous theorem, and define $\phi_{k,n} = \int\int_{\mathcal{R}_{k,n}} -\kappa dA + \int_{r(I_k)} \mu_{\mathcal{R}_{k,n}}$.

Then as we saw in the previous theorem, Theorem 8.7, there is $K_0 > 0$ such that

$$\phi_{k,n} \geq K_0 \tag{8.6}$$

We continue to show that after possibly passing to a subsequence there are parameter values $\hat{t}_n$ for which $g_n(\hat{t}_n)$ converge to a point $z \in \mathrm{Teich}(S)$.

Let $T > 0$ be such that $d(r(t), r'(t)) > \delta$ for every $t > T$. We claim that there is $\hat{k}$ such that $\mathcal{R}_{\hat{k},n} \cap g_n \neq \emptyset$ for every sufficiently large n.

Assume that $g_n \cap \mathcal{R}_{k,n} = \emptyset$ for every $k \leq N$ then $\mathcal{R}_{k,n} \subset \mathfrak{R}_n$. Thus by (8.6) we get the following estimate

$$\int\int_{\mathfrak{R}_n} -\kappa dA + \int_{r([T,n])} \mu_{\mathfrak{R}_n} > NK_0$$

Further by the Gauss-Bonnet theorem the left hand side is bounded above by π independent from N, so we get the upper bound $\frac{\pi}{K_0}$ for N.

Now if there is no $\hat{k}$ such that all g_n intersect $\mathcal{R}_{\hat{k},n}$, then N can be made arbitrary large and we get a contradiction to the upper bound above and the claim follows.

Let $z_n = g_n(\hat{t}_n)$ be the intersection points. Then z_n lie in the s tubular neighborhood of $r(I_{\hat{k}})$ which is a compact subset of $\mathrm{Teich}(S)$, therefore after passing to a subsequence we may assume that z_n converge to a point $z \in \mathrm{Teich}(S)$.

Now we proceed to show that the geodesic segments g_n converge to a geodesic passing through z. We claim that if we reparametrize g_n by arc-length such that $g_n(0) = z_n$ then for each $t \in [0, \infty)$ the sequence $\{g_n(t)\}_n$ is Cauchy. To see this let h_n^+ be the geodesic segment joining z to $r(n)$ parametrized by arc-length, then choose N_t such that $d(z, r(n)) > t$ for each $n \geq N_t$. Now $d(h_n^+(t), g_n^+(t)) < d(z_n, z) \to 0$ as $n \to \infty$ thus the claim follows from a CAT(0) comparison for the triangles $\triangle(zz_nr(n))$.

A similar argument shows that for each $t \in (-\infty, 0]$ $\{g_n(-t)\}_n$ is Cauchy.

Therefore g_n limit to a piece-wise geodesic $g : \mathbb{R} \to \overline{\mathrm{Teich}(S)}$ as parametrized geodesics. Moreover by Proposition 8.2 in [BH99] II.8 g is a bi-infinite geodesic in $\overline{\mathrm{Teich}(S)}$.

Moreover by the construction $g^+ = g|_{[0,\infty)}$ is asymptotic to r and $g^- = g|_{[0,-\infty)}$ is asymptotic of r'. Also as we saw in Theorem 8.7 g^+ is actually strongly asymptotic to r and the lemma follows.

□

Proposition 8.10. *Let r_ρ be a ray with prescribed itinerary with narrow ending data (ν^-, ν^+) and bounded annular coefficients. Let μ be a measure supported on ν^+. Then the length-function $\ell_\mu(r_\rho(t))$ is bounded.*

Proof. As Lemma 4.5 in [BMM10] we consider the ray $\hat{r}$ as follows:

By the structure of hierarchy paths $\nu^+ = \nu' \in \mathcal{EL}(Z)$ where $Z \subseteq S$. Let $\Sigma_{\nu'}$ denote the simplex of projective classes of measures on ν' in $\mathcal{PML}(Z)$, and let $\hat{\mu} \in \mathcal{ML}(Z)$ be a representative of the projective class determined by a point in the interior of the top-dimensional face. Then $\hat{\mu}$ is a positive linear combination of all ergodic measures on ν'. Let γ_n be a sequence of simple closed curves for which the projective classes $[\gamma_n]$ converge to $[\hat{\mu}]$, and let $r_n = [x, z_n]$ where $z_n \in \mathcal{S}(\gamma_n)$. Since γ_n are pinching curves for r_n, Lemma 3.14 implies that any limit $\hat{r}$ of a convergent subsequence of r_n has the property that $\hat{\mu}$

has bounded length along $\hat{r}$. Since $\hat{\mu}$ is a positive linear combination of all the ergodic measures on ν^+, it follows that each ergodic measure on ν^+ has bounded length along $\hat{r}$. Hence, any measured lamination representing a projective class in Σ has bounded length along $\hat{r}$ since each is a linear combination of ergodic measures.

$\hat{r}$ is an infinite ray in $\mathcal{S}(\partial Z)$, since otherwise $\hat{r}$ would have a pinching curve γ with $\iota(\gamma, \hat{\mu}) > 0$, violating the length bound on $\hat{\mu}$ along $\hat{r}$.

Let $\bar{\mu}$ be any ending measure for r_ρ then $\bar{\mu}$ decomposes to a measure with a projective representative in Σ and a measure on ∂Z, thus has bounded length along $\hat{r}$.

Now observe that r_ρ and $\hat{r}$ are strongly asymptotic to each other, since otherwise Theorem 8.9 implies that we have a bi-infinite geodesic g asymptotic to r_ρ in forward time and asymptotic to $\hat{r}$ in backward time along which $\bar{\mu}$ has bounded length, which contradicts the convexity of the length-function of $\bar{\mu}$ along g.

Now μ decomposes to a measure μ' supported on ν' which is represented in Σ and a measure supported on ∂Z. Since r_ρ and $\hat{r}$ are strongly asymptotic μ' has bounded length along r_ρ, moreover by Corollary 8.3

$$\ell_{\partial Z}(r_\rho(t)) \to 0$$

as $t \to \infty$, thus the component supported on ∂Z also has bounded length along r_ρ. Therefore μ has bounded length along r_ρ and the proposition follows. $\square$

Proof of Theorem 1.3. Assume that r and r' are not strongly asymptotic. Then by Theorem 8.9 there is a bi-infinite geodesic g strongly asymptotic to r in forward time and asymptotic to r' in backward time.

Let μ be an ending measure of r' obtained as a weighted limit of Bers curves at $\{r'(n)\}_n$. Then since g_n has left end point $r'(n)$, μ is an ending measure of $g^- = g|_{[0,-\infty)}$ as well. Therefore by Lemma 3.13 it has bounded length in backward time along g.

Moreover, since the forward ending lamination of r' is ν^+, μ is supported on ν^+, thus as we saw in Proposition 8.10 μ has bounded length along r as well.

Now since r and $g^+ = g|_{[0,\infty)}$ are strong asymptotic by the continuity of length-functions μ has bounded length along g^+.

Therefore, the measured lamination μ has bounded length both backward and forward along g which contradicts the convexity of the $\mu-$length-function along WP geodesics. The theorem follows from this contradiction.

$\square$

9 Appendix : Ending data with prescribed subsurface coefficients

Our purpose in this appendix is to provide ending data, pairs of partial markings or laminations, with a given list of large subsurface coefficients. We borrow several ideas and notations from [CLM12], [Sun].

We start by recalling the following two results about the action of psuedo-Anosov maps on the curve complex of a surface and its subsurfaces.

Theorem 9.1. *[MM00]*
Let f be a pseudo-Anosov map supported on $Y \subseteq S$, then there is a constant $m(f)$ depending on f, such that for any subsurface $W \neq Y$ and any partial marking μ we have:

$$\max\{d_W(\mu, f^e\mu) : e \in \mathbb{Z}\} \leq m(f)$$

Theorem 9.2. *[MM99]*
Let f be a pseudo-Anosov map supported on $Y \subseteq S$, then there is the positive number $\tau_Y(f)$, called the translation length of f such that for every partial marking μ and $e \in \mathbb{Z}$ we have:

$$d_Y(\mu, f^e\mu) > \tau_Y(f)|e|$$

Let $\{X_i\}_{i=1}^n$ be any finite collection of subsurfaces of S including annuli, also assume that any two of them are either nested or overlap. For each i, let f_i be a pseudo-Anosov map supported on X_i. We continue to set up some definitions and notations:

Let $q : \{1, 2, ..., k\} \to \{1, 2, ..., n\}$ and $e : \{1, 2, ..., k\} \to \mathbb{Z}$, we set up the following sequence of subsurfaces of S, for each $i = 1, ..., k$ let

$$X^{q,e}(f_{q(i)}) = f_{q(1)}^{e_1} ... f_{q(i-1)}^{e_{i-1}} X_{q(i)}$$

Moreover, for each i let j be the largest index such that $j < i$ and $X^{q,e}(f_{q(j)}) \pitchfork X^{q,e}(f_{q(i)})$, and define $\mu_{q,e}^-(i) = \partial X^{q,e}(f_{q(j)})$.

Similarly, let j be the smallest index such that $j > i$ and $X^{q,e}(f_{q(j)}) \pitchfork X^{q,e}(f_{q(i)})$, and define $\mu_{q,e}^+(i) = \partial X^{q,e}(f_{q(j)})$.

Let $I_{q,e} = \{i : \mu_{q,e}^-(i) = \emptyset\}$ and let $\mu_{q,e}^-$ be any partial marking containing $\{\partial X^{q,e}(f_{q(i)}) : i \in I_{q,e}\}$. Similarly, let $T_{q,e} = \{i : \mu_{q,e}^+(i) = \emptyset\}$ and let $\mu_{q,e}^+$ be any partial marking containing $\{\partial X^{q,e}(f_{q(i)}) : i \in T_{q,e}\}$.

For any i if both $\mu_{q,e}^-(i)$ and $\mu_{q,e}^+(i)$ are non empty then we call i an interior index.

Now we are ready to state the following lemma of Brock, Masur and Minsky. This lemma and its generalization appear in [Sun].

Lemma 9.3. *(Local to Global lemma) There are constants M_0 and L_0 depending only on the topological type of S with the following property: Given q, e let $Z_i = X^{q,e}(f_{q(i)})$. Then for an interior index i if $d_{Z_i}(\mu_{q,e}^-(i), \mu_{q,e}^+(i)) > M_0$ then*

$$d_{Z_i}(\mu_{q,e}^-, \mu_{q,e}^+) \asymp_{1,L_0} d_{Z_i}(\mu_{q,e}^-(i), \mu_{q,e}^+(i))$$

The following propositions are consequences of the Local to global lemma. In the rest of this appendix we assume that q and e are such that the collection of surfaces $\{X^{q,e}(f_{q(i)})\}_{i=1}^k$ fills the surface S, that is every subsurface $Y \subseteq S$ overlap at least one subsurface in the collection, also let $M = \max\{M_1, M_2, B_0\}$.

Proposition 9.4. *(Large domains) Given* q *and* e *let* $Z_i = X^{q,e}(f_{q(i)})$*. There is a* $L \geq 0$ *such that for any interior index* i *we have*

$$d_{Z_i}(\mu^-_{q,e}, \mu^+_{q,e}) \asymp_{1,L} \bar{\tau}|e_i| \tag{9.1}$$

where $\bar{\tau} = \min_{i=1,\dots,n} \tau_{X_i}(f_i)$*.*

Proof. First we establish the lower bound in (9.1).

Claim 9.5. There is $L' > 0$ for which $d_{Z_i}(\mu^-_{q,e}(i), \mu^+_{q,e}(i)) \geq \bar{\tau}|e_i| - L'$

Note that if the claim holds and if $|e_i|$ be large enough such that $\bar{\tau}|e_i| - L_0 > M_0$ then the local to global lemma applies and we get

$$d_{Z_i}(\mu^-_{q,e}, \mu^+_{q,e}) > d_{Z_i}(\mu^-_{q,e}(i), \mu^+_{q,e}(i)) - L' - L_0 > \bar{\tau}|e_i| - L' - L_0$$

which gives the lower bound in (9.1).

By the definition of $\mu^{\pm}_{q,e}(i)$ there are j and j' such that $\partial Z_j = \mu^-_{q,e}(i)$ and $\partial Z_{j'} = \mu^+_{q,e}(i)$. Let $a = f^{e_1}_{q(1)} \dots f^{e_{i-1}}_{q(i-1)}$ applying a^{-1} we get

$$d_{Z_i}(\partial Z_j, \partial Z_{j'}) = d_{X_{q(i)}}(c\partial X_{q(j)}, f^{e_i}_{q(i)} c' \partial X_{q(j')}) \tag{9.2}$$

in the above expression $c = \prod_{j<i'<i} f^{-e_{i'}}_{q(i')}$ and $c' = \prod_{i<i'<j'} f^{e_{i'}}_{q(i')}$. Now the triangle inequality in $\mathcal{C}(X_{q(i)})$ implies that the right hand side of (9.2) is bounded below by

$$d_{X_{q(i)}}(f^{e_i}_{q(i)} c' \partial X_{q(j)}, c' \partial X_{q(j')}) - d_{X_{q(i)}}(c' \partial X_{q(j')}, c \partial X_{q(j)}) \tag{9.3}$$

We claim that the second term in (9.3) is bounded above by $\xi(S)\bar{m}$, where $\bar{m} = \max_{i=1,\dots,n} m(f_i)$. By the triangle inequality it is bounded above by

$$d_{X_{q(i)}}(f^{e_{i'}}_{q(i')} c'' \partial X_{q(j')}, c'' \partial X_{q(j')}) + d_{X_{q(i)}}(c'' \partial X_{q(j')}, c \partial X_{q(j)})$$

where $c'' = f^{-e_{i'}}_{q(i')} c'$, now since $X_{q(i')} = \mathrm{supp}(f_{q(i')}) \neq X_{q(i)}$ Theorem 9.1 implies that the first term above is bounded above by $\bar{m}$, so we are left with the term

$$d_{X_{q(i)}}(c'' \partial X_{q(j')}, c \partial X_{q(j)})$$

Now by repeating the procedure we used to bound from above the second term in (9.3) noting the fact that for each i', $\mathrm{supp}(f_{q(i')}) \neq X_{q(i)}$ our claim follows.

Moreover, by Theorem 9.1 the first term in (9.3) is larger than $\tau_{X_{q(i)}}(f_{q(i)})$ thus for $L' = \xi(S)\bar{m}$ Claim 9.5 follows.

To get the upper bound in (9.1) note that the right hand side of (9.2) is bounded above by

$$d_{X_{q(i)}}(f^{e_i}_{q(i)} \partial X_{q(j)}, c' \partial X_{q(j')}) + d_{X_{q(i)}}(c' \partial X_{q(j')}, c \partial X_{q(j)})$$

then the exact same argument as before gives us the bound $\bar{\tau}|e_i|$ for the first term and L' for the second term in the above expression and the upper bound follows.

$\square$

Proposition 9.6. *(Order of domains) Given* q *and* e*, let* $Z_i = X^{q,e}(f_{q(i)})$*. If* $i < j$ *and* $Z_i \pitchfork Z_j$ *then* $Z_i < Z_j$ *in any hierarchy path between* $\mu^-_{q,e}$ *and* $\mu^+_{q,e}$*.*

Proof. Let $q_{int} = q|_{\{1,\dots,i\}}$ and $e_{int} = e|_{\{1,\dots,i\}}$ then $\mu^-_{q_{int},e_{int}} \subset \mu^-_{q,e}$ and $\partial Z_j \subset \mu^+_{q_{int},e_{int}}$. Thus Proposition 9.4 for q_{int} and e_{int} implies that

$$d_{Z_i}(\mu^-_{q,e}, \partial Z_j) \asymp_{1,L} \bar{\tau}|e_i|$$

so for $|e_i|$ sufficiently large we get $d_{Z_j}(\partial Z_i, \mu^-_{q,e}) \geq M$.

Similarly, considering $q_{ter} = q|_{\{j,\dots,k\}}$ and $e_{ter} = e|_{\{j,\dots,k\}}$ we get $d_{Z_i}(\partial Z_j, \mu^+_{q,e}) \geq M$.

The above two inequalities as we saw in Proposition 2.14 imply that $Z_i < Z_j$ in any hierarchy path between $\mu^-_{q,e}$ and $\mu^+_{q,e}$.

$\square$

Proposition 9.7. *Given* q *and* e*, let* $Z_i = X^{q,e}(f_{q(i)})$*. There is a constant* $m > M_1$ *only depending on the pseudo-Anosov maps* $\{f_i\}_{i=1}^n$ *such that for any subsurface* $Z \notin \{Z_i\}_{i=1}^k$

$$d_Z(\mu^-_{q,e}, \mu^+_{q,e}) \leq m$$

Proof. Let $\bar{m} = \max_{i=1,\dots,n} m(f_i)$. By our assumption $\{Z_i\}_i$ is filling, thus there is at least one index l such that $Z_l \pitchfork Z$. If $d_Z(\mu^-_{q,e}, \mu^+_{q,e}) \leq M_1$ we already have the bound, otherwise Z is a component domain of any hierarchy path between $\mu^-_{q,e}$ and $\mu^+_{q,e}$. Moreover by Proposition 9.4 each Z_i is a component domain of any such hierarchy path. In the case that there are at least two indices l and l' such that $Z_l < Z < Z_{l'}$ pick two indices with $l' - l \leq \xi(S)$ then by Theorem 2.11 (5) we have

$$d_Z(\mu^-_{q,e}, \mu^+_{q,e}) \leq d_Z(\partial Z_l, \partial Z_{l'}) + 2M_2$$

let $a = f^{e_1}_{q(1)} \dots f^{e_l}_{q(l)}$, applying a^{-1} to the right hand side of the above inequality we get

$$d_{a^{-1}Z}(\partial X_{q(l)}, c\partial X_{q(l')}) + 2M_2$$

where $c = f^{e_l}_{q(l)} \dots f^{e_{l'-1}}_{q(l'-1)}$.

We claim that the first term above is bounded above by $\xi(S)\bar{m}$, to see this note that $a^{-1}Z \neq \mathrm{supp}(f_{q(j)})$ for any $l < j < l'$, thus the claim follows by applying Theorem 9.1 repeatedly.

In the case that there is only one index l either $Z < Z_l$ and $l \leq \xi(S)$ or $Z_l < Z$, in the first case we have

$$d_Z(\mu^-_{q,e},\mu^+_{q,e}) \le d_Z(\mu^-_{q,e},\partial Z_l) + 2M_2$$

but then Theorem 9.1 applies directly and implies that the first term above is bounded above by $\xi(S)\bar{M}$, because $Z \ne \mathrm{supp}(f_{q(j)})$ for any $1 < j < l$. The second case can be treated exact the same way by considering q' and e' defined respectively by $q'(i) = q(k-i+1)$ and $e'(i) = e(k-i+1)$.

□

Now we can put the above three propositions together and give a recipe for a hierarchy with a prescribed list of large domains.

Start with subsurfaces $\{X_i\}_{i=1}^n$, pseudo-Anosov maps $\{f_i\}_{i=1}^n$ such that for each i, f_i is supported on X_i, and functions $q:\{1,2,...,k\}\to\{1,2,..,n\}$ and $e:\{1,2,...,k\}\to\mathbb{Z}$.

Recall subsurfaces $Z_i = f^{e_1}_{q(1)}...f^{e_{i-1}}_{q(i-1)}X_{q(i)}$, and let $\bar{\tau}$ and L be as in Proposition 9.4 and m be as in Proposition 9.7, then the two markings $\mu^-_{q,e}$ and $\mu^+_{q,e}$ have the following properties:

By Proposition 9.4 we have

$$d_{Z_i}(\mu^-_{q,e},\mu^+_{q,e}) \asymp_{1,L} \bar{\tau}|e_i|$$

moreover, by Proposition 9.7 for any $Z \notin \{Z_i\}_i$ we have

$$d_{Z_i}(\mu^-_{q,e},\mu^+_{q,e}) \le m$$

The above bounds prescribe all the subsurface coefficients of the pair $(\mu^-_{q,e},\mu^+_{q,e})$ and consequently all the *large domains* of any hierarchy path with ending data $(\mu^-_{q,e},\mu^+_{q,e})$.

Moreover, Proposition 9.6 prescribes the partial order of Z_i's in any hierarchy path with ending data $(\mu^-_{q,e},\mu^+_{q,e})$.

To generalize this construction to infinite hierarchies, let $q:\mathbb{N}\to\{1,...,n\}$ and $e:\mathbb{N}\to\mathbb{Z}$ be any sequence of integers. Now consider hierarchies paths with ending data (μ,μ_k) as above, where $\mu \equiv \mu^-_{q_k,e_k}$ and $\mu_k = \mu^+_{q_k,e_k}$ for the function $q_k = q|_{\{1,2,...,k\}}$ (which is the restriction of q to $\{1,2,...,k\}$) and $e_k = e|_{\{1,2,...,k\}}$. Then using what we have shown in the finite case we have:

(i) Given i then $d_{Z_i}(\mu,\mu_k) \asymp_{1,L} \bar{\tau}n_i$ for every $k > i$, and

(ii) If $Z \notin \{Z_i\}_i$ then $d_Z(\mu,\mu_k) \le m$ for every k.

After possibly passing to a subsequence $[\mu_k] \to [\mu_\infty]$ in $\mathcal{PML}(S)$. Therefore, $|\mu_k| \to |\mu_\infty| \cup \{\text{finitely many isolated leaves}\}$ in the Hausdorff topology of closed subsets of S, see for example [CEG87]. The Hausdorff convergence implies that $d_{Z_i}(\mu,\mu_\infty) \asymp_{1,1} d_{Z_i}(\mu,\mu_k)$.

By the above bounds for each subsurface Z_i and k sufficiently large we have

$$d_{Z_i}(\mu,\mu_\infty) \asymp_{1,1} d_{Z_i}(\mu,\mu_k) \asymp_{1,L} \bar{\tau}n_i$$

moreover, for every $Z \notin \{Z_i\}_i$

$$d_Z(\mu, \mu_\infty) = d_Z(\mu, \mu_k) \leq m$$

also order of domains is the same as the order of them in the finite case, that is if $i < j$ and $Z_i \pitchfork Z_j$ then $Z_i < Z_j$ in any hierarchy path with ending data (μ, μ_∞).

References

[BDM09] Jason Behrstock, Cornelia Druţu, and Lee Mosher, *Thick metric spaces, relative hyperbolicity, and quasi-isometric rigidity*, Math. Ann. **344** (2009), no. 3, 543–595.

[Beh06] Jason A. Behrstock, *Asymptotic geometry of the mapping class group and Teichmüller space*, Geom. Topol. **10** (2006), 1523–1578.

[BF06] Jeffrey Brock and Benson Farb, *Curvature and rank of Teichmüller space*, Amer. J. Math. **128** (2006), no. 1, 1–22.

[BH99] Martin R. Bridson and André Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften, vol. 319, Springer-Verlag, Berlin, 1999.

[BKMM12] Jason Behrstock, Bruce Kleiner, Yair Minsky, and Lee Mosher, *Geometry and rigidity of mapping class groups*, Geom. Topol. **16** (2012), no. 2, 781–888.

[BM08] Jeffrey Brock and Howard Masur, *Coarse and synthetic Weil-Petersson geometry: quasi-flats, geodesics and relative hyperbolicity*, Geom. Topol. **12** (2008), no. 4, 2453–2495.

[BMM10] Jeffrey Brock, Howard Masur, and Yair Minsky, *Asymptotics of Weil-Petersson geodesics. I. Ending laminations, recurrence, and flows*, Geom. Funct. Anal. **19** (2010), no. 5, 1229–1257.

[BMM11] ______, *Asymptotics of Weil-Petersson geodesics II: bounded geometry and unbounded entropy*, Geom. Funct. Anal. **21** (2011), no. 4, 820–850.

[Bro] Jeffrey Brock, *Chains of flats and non-unique ergodicity of Weil-Petersson ending laminations*, In preparation.

[Bro03] Jeffrey F. Brock, *The Weil-Petersson metric and volumes of 3-dimensional hyperbolic convex cores*, J. Amer. Math. Soc. **16** (2003), no. 3, 495–535.

[Bus10] Peter Buser, *Geometry and spectra of compact Riemann surfaces*, Modern Birkhäuser Classics, Birkhäuser Boston Inc., Boston, MA, 2010, Reprint of the 1992 edition.

[CE08] Jeff Cheeger and David G. Ebin, *Comparison theorems in Riemannian geometry*, AMS Chelsea Publishing, Providence, RI, 2008, Revised reprint of the 1975 original.

[CEG87] R. D. Canary, D. B. A. Epstein, and P. Green, *Notes on notes of Thurston*, Analytical and geometric aspects of hyperbolic space (Coventry/Durham, 1984), London Math. Soc. Lecture Note Ser., vol. 111, Cambridge Univ. Press, Cambridge, 1987, pp. 3–92.

[CLM12] Matt T. Clay, Christopher J. Leininger, and Johanna Mangahas, *The geometry of right-angled Artin subgroups of mapping class groups*, Groups Geom. Dyn. **6** (2012), no. 2, 249–278.

[CM06] Yitwah Cheung and Howard Masur, *A divergent Teichmüller geodesic with uniquely ergodic vertical foliation*, Israel J. Math. **152** (2006), 1–15.

[DW03] Georgios Daskalopoulos and Richard Wentworth, *Classification of Weil-Petersson isometries*, Amer. J. Math. **125** (2003), no. 4, 941–975.

[Ebe72] Patrick Eberlein, *Geodesic flows on negatively curved manifolds. I*, Ann. of Math. (2) **95** (1972), 492–510.

[Eck93] Jürgen Eckhoff, *Helly, Radon, and Carathéodory type theorems*, Handbook of convex geometry, Vol. A, B, North-Holland, Amsterdam, 1993, pp. 389–448.

[Ham] Ursula Hamenstädt, *Closed Teichmüeller geodesics in the thin part of moduli space*, arxiv.

[Kla] Erica Klarreich, *The boundary at infinity of the curve complex*, Preprint, 1999.

[Kli78] Wilhelm Klingenberg, *A course in differential geometry*, Springer-Verlag, New York, 1978, Translated from the German by David Hoffman, Graduate Texts in Mathematics, Vol. 51.

[Mah10] Joseph Maher, *Linear progress in the complex of curves*, Trans. Amer. Math. Soc. **362** (2010), no. 6, 2963–2991.

[Mas76] Howard Masur, *Extension of the Weil-Petersson metric to the boundary of Teichmuller space*, Duke Math. J. **43** (1976), no. 3, 623–635.

[Mas09] ———, *Geometry of Teichmüller space with the Teichmüller metric*, Surveys in differential geometry. Vol. XIV. Geometry of Riemann surfaces and their moduli spaces, Surv. Differ. Geom., vol. 14, Int. Press, Somerville, MA, 2009, pp. 295–313.

[McM00] Curtis T. McMullen, *The moduli space of Riemann surfaces is Kähler hyperbolic*, Ann. of Math. (2) **151** (2000), no. 1, 327–357.

[Min01] Yair N. Minsky, *Bounded geometry for Kleinian groups*, Invent. Math. **146** (2001), no. 1, 143–192.

[Min10] Yair Minsky, *The classification of Kleinian surface groups. I. Models and bounds*, Ann. of Math. (2) **171** (2010), no. 1, 1–107.

[MM99] Howard A. Masur and Yair N. Minsky, *Geometry of the complex of curves. I. Hyperbolicity*, Invent. Math. **138** (1999), no. 1, 103–149.

[MM00] H. A. Masur and Y. N. Minsky, *Geometry of the complex of curves. II. Hierarchical structure*, Geom. Funct. Anal. **10** (2000), no. 4, 902–974.

[Mun75] James R. Munkres, *Topology: a first course*, Prentice-Hall Inc., Englewood Cliffs, N.J., 1975.

[PP10] Jouni Parkkonen and Frédéric Paulin, *Prescribing the behaviour of geodesics in negative curvature*, Geom. Topol. **14** (2010), no. 1, 277–392.

[PWW10] Mark Pollicott, Howard Weiss, and Scott A. Wolpert, *Topological dynamics of the Weil-Petersson geodesic flow*, Adv. Math. **223** (2010), no. 4, 1225–1235.

[Raf] Kasra Rafi, *Closed geodesics in the thin part of moduli space*, In preparation.

[Raf05] ______, *A characterization of short curves of a Teichmüller geodesic*, Geom. Topol. **9** (2005), 179–202.

[Raf07] ______, *A combinatorial model for the Teichmüller metric*, Geom. Funct. Anal. **17** (2007), no. 3, 936–959.

[Ser85] Caroline Series, *The modular surface and continued fractions*, J. London Math. Soc. (2) **31** (1985), no. 1, 69–80.

[Sun] Chun-yi Sun, *Asymptotic geometry, bounded generation and subgroups of mapping class groups*, PhD thesis, Yale University, 2012.

[Wol03] Scott A. Wolpert, *Geometry of the Weil-Petersson completion of Teichmüller space*, Surveys in differential geometry, Vol. VIII (Boston, MA, 2002), Surv. Differ. Geom., VIII, Int. Press, Somerville, MA, 2003, pp. 357–393.

[Wol08] ______, *Behavior of geodesic-length functions on Teichmüller space*, J. Differential Geom. **79** (2008), no. 2, 277–334.

[Wol10] ———, *Families of Riemann surfaces and Weil-Petersson geometry*, CBMS Regional Conference Series in Mathematics, vol. 113, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2010.

[Wu] Yunhui Wu, PhD thesis, Brown University, 2012.

[Yam04] Sumio Yamada, *On the geometry of Weil-Petersson completion of Teichmüller spaces*, Math. Res. Lett. **11** (2004), no. 2-3, 327–344.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 442 DUNHAM LAB., 10 HILLHOUSE AVE., NEW HAVEN, CT, 06511
E-mail address: `babak.modami@yale.edu`